

## **BIBLIOGRAPHY**

---

## BIBLIOGRAPHY

- A.O.A.C. (1970). Official Methods of Analysis, (XI Edn.), Washington D.C.
- A.O.A.C. (1975). Official Methods of Analysis, (XII Edn.), Washington D.C.
- Abraham Z. (1981). Glimpses of Indian ethno-botany. Oxford and IBH publishing Co., New Delhi, pp. 308
- Ahmed, S.A. (2012). Exploration of forest bio-resources in vanya silk sector with special reference to eri silk. *MUCCALUNG*. International Consultative Meeting on Seri-biotechnology 2012 (ICMS 2012), December 5-7, 2012, Imphal, Manipur (India). pp.20
- Ahmed, S.A. and Rajan, R.K. (2011). Exploration of *vanya* silk biodiversity in North Eastern Region of India: Sustainable livelihood and poverty alleviation. Proceedings of International Conference on Management, Economics and Social Sciences (ICMESS'2011), December 23-24, 2011 at Bangkok, Thailand. pp. 485-489
- Ahmed, S.A., Goswami, D and Mech, D. (2012). Effect of pesticide applied in tea gardens on muga silkworm rearing. *Compendium of Seminar Papers*. National Seminar on Recent Trends in Research & Development in Muga Culture: Ideas to Action, 3-4 May 2012, Guwahati, pp: 114-121.
- Arora, G.S. and Gupta, I.J. (1979). Taxonomic studies of some of the Indian non-mulberry silk moths (Lepidoptera: Saturniidae). *Mem. Zool.Surv. India.*, 16(1): 1-64.
- Ataa Said; Usama W. Hawas; Siham El-Shenawy; Salwa M. Nofal and Khaled Rashed (2010). Flavonoids and some biological activities of *Ailanthus excelsa* Leaves, *IUFS J Biol* 2010, 69(1): 41-55
- Baker, J. E. and Mabie, J. A. (1973). Feeding behaviour of larvae of *Plodia interpunctella*. *J. Econ. Entomol.* 2: 627-632

---

Banerjee, N.D.; Choudhury, A.K.; Sinha, U.S.P. and Brahmachari, B.N. (1993). Studies on the foliar constituents of the food plants of temperate tasar silkworm, *Anthearea proylei* Jolly., *Indian J. Seric.*, **32**(2): 228-230.

Barah, A.; Goswami, M.C. and Samson, M.V. (1989). Consumption and utilization of food in different instars of muga silkworm *Antheraea assama* Westwood, *Proc. Indian Acad. Sci. (Anim. Sci.)*, **98**(2): 99-103.

Barthlott, William, Linsenmair, K. Eduard and Porembski, Stefan (2005). Biodiversity: Structure and Function. Vol. I, Encyclopedia of Life Support System.

Basak, R. K. (1980). in *Fascicles of Flora of India*, Fascicle 4, pp 1-7. Botanical Survey of India, Botanic Garden.

Bernays E.A. and Chapman, R.F. (1994). Host-Plant Selection by Phytophagous Insects. Chapman and Hall, London, pp. 61-94.

Bhattacharya, A.K. and Pant, N.C. (1976). Studies on the insect host plant relationship. Consumption and utilization profile in insect. *Proc. Nat. Acad. Sci. India*, **46** (8): 273-299

Bindu P.C., Jaisankar P., Hauer F., Gutzeit H.O. and Kundu, S.C. (2006). Biological relevance of host plant derived terpinoid in the cocoons of the tropical tasar silkworm *Antheraea mylitta*, *Biochemical Systematics and Ecology*, **34**(9): 698-704.

Bongale U. D. and Chaluvachari K. (1993) Evaluation of eight mulberry varieties by leaf biochemical and bioassay moulting studies. *Sericologia*. **35** (1): 83-94.

Bose, P.C.; Majumdar, S.K. and Sengupta, K. (1991). A comparative biochemical study of six mulberry (*Morus alba* L.) varieties. *Indian J. Seric.*, **30**(1): 83-87.

British Pharmacopoeia (1988). II, H. M. Stationary office: London, 704.

Brues, C.T. (1946). Insect Dietary. Cambridge, Mass. pp.466

---

Buddhi Katangodage (2009). Food Science Experiment- Crude Fat Analysis. Link:  
<http://experiments-budrockz.blogspot.in/2009/09/food-science-experiment-crude-fat.html>

Cantelo. W. W. and M. Jacobson (1979). Phenacetaldehyde attracts moths to bladder flower and to black light traps. *Environmental Entomology*, 8: 444-447.

Chakravorty, R.; Neog, K.; Suryanarayana, N. and Hazarika, L.K. (2004). Feeding and moulting behaviour of muga silkworm (*Antheraea assama*) on different food plants. *Sericologia*, 44(2): 145-152.

Chaluvachari, K. and Bongale, U.D. (1995). Leaf quality evaluation of selected mulberry genotypes by biochemical and bioassay studies, *Proceedings of Conference on Cytology and Genetics*, 4: 121-124.

Chandrashekhar S.; Sannappa B.; Manjunath K.G. and Govindan R. (2013). Nutritive value of leaves in different genotypes of castor (*Ricinus communis* L.), *Indian Journal of Plant Sciences*, 2 (2): 22-27

Chaturvedi, M D. (1956). *Ardu*, the tree of distinction, *Ind Fmg*, 5(1956):32-34.

Chopra R.N.; Nayar S.L. and Chopra I.C. (1956). Glossary of Indian medicinal plants. CSIR, New Delhi, p 10.

Choudhury, S.N.; Deka, D.K. and Rao, P.G. (2006). Role of biochemical constituents of som plant (*Persea bombycina*) on muga crop production, Proceedings of workshop on Problems and Prospects of Muga Silkworm Seed Production. Pp. 46-50.

Chowdhury, S.N. (1968). *Indian Silk*, published by Central Silk Board, 6(1968):9

Chowdhury, S.N. (1982). Eri Silk Industry, published by Dir Seri. & Wvg, Guwahati.

Chowdhury, S.N. (2004). Origin, evolution and distribution of silkworm races, *J Assam Sc Soc*, 45: 43-51.

---

Chowdhury, S.N. (2006). Host Plant species of Non-mulberry (Vanya) Silkworms. Non-Mulberry Silkworm and Host Plants Germplasm. pp.67-73

Clarke, T.M. and Townby, S.T.F. (1986). Heritability and relationship to yield of excised leaf water retention in durum wheat. *Crop Sci.*, 26(2): 289-292.

Dastur J.F. (1951). Medicinal plants of India & Pakistan, 1st edn. D.B. Taraparevala Sons and Co., Bombay, p 18

Dethier, V. G.; Barton, B. L. and Smith, C. N. (1960). The designation of chemicals in terms of the responses they elicit from insects. *J. Econ. Entomol.* 53: 134-136.

Devaiah, M. C.; Rajashekargouda, R. and Suhas Yelshetty (1988). Effect of castor, tapioca and their combinations on the growth and development of the larvae and cocoon characters of *Samia cynthia ricini* Boisduval (Abstract). *International Congress on Tropical Sericulture Practices, Bangalore* February, 1988, p. 33.

Dhanasekaran S.; Suresh B.; Sethuraman M.; Rajan S. and Dubey R. (1993). Antifertility activity of *Ailanthus excelsa* Linn. In female albino rats. *Indian Journal of Experimental Biology* 31, 384-385.

Dookia, B.R. (1980). Varied silk ratio in cocoons of eri silkworm (*Philosamia ricini* Hutt.) reared on different castor varieties in Rajasthan. *Indian Journal of Sericulture*, 19: 38-40.

Dutta, L.C. (2000). Effect of castor varieties on growth, nutrition and cocoon characters of eri silkworm, *Samia cynthia ricini* Boisduval, Ph.D. Thesis, AAU, Jorhat.

Dutta, L.C. and Kalita, D.N. (1997). Food consumption and utilization by the larvae of Eri silk (*Philosamia ricini* Hutt.) in different food plants. *J. Appl. Zool. Res.* 8(2): 143-144.

---

Dutta, L.C.; Kalita, M.N. and Sarkar, C.R. (1997). Foliar constituents of the food plants of muga silkworm *Antheraea assama* Westwood. *Indian J. Seric.* **36** (1): 85-86.

Edreva A.; Velikova V.; Tsonev T.; Dagnon S.; Gürel A.; Aktaş L. and Gesheva E. (2008). Stress protective role of secondary metabolites: Diversity of functions and mechanisms, *Gen. Appl. Plant Physiol.*, Special Issue, 34 (1-2): 67-78.

Farlane Mc JE, and Distler M.H.W., The effect of rutin on growth, fecundity and food utilization in *Achilus domesticus* L., *J. Insect Physiol.*, 28: 85-88.

Fonseca T. C.; De Almeida J. E.; Okamoto F. and Da Cunha E. A. (1993). Nutritive value of some mulberry clones (*Morus alba* L.). *Sericologia*. **33**(4): 631-633.

Foster, S. P. and Harris, M. O. (1997). Behavioural manipulation methods for insect pest management. *Annu. Rev. Entomol.* **42**: 123-146.

Fotedar, R.K. and Dandin, S.B. (1997). Chemical composition and feeding studies of different elite mulberry genotype under temperate conditions, *Indian J. Seric.* **36** (1): 22-26.

Fraenkel, G.S. (1959). The raison d'etre of secondary plant substances. *Science*. **129**: 1466-70.

Fukuda T.; Suda M.; Matsuda M.; Mayashi T.; Kuroso T.; Horiuchi Y. and Florrin M. (1959) Formation of the silk protein during the growth of the silkworm. *Biochemistry*. **12**: 90-112.

Fukuda, T. (1960). The correlation between the mulberry leaves taken by the silkworm, the silk protein in the silk gland and the silk filament, *Bull. Seric. Exp. Stn.* **15**(4): 605-610.

Fukuda, T.; Kamegama, T. And Matsuda, M. (1963). A correlation between the mulberry leaves consumed by the silkworm larva in different ages of the larval growth

---

- 
- and production of the fibre spun by the silkworm larva and the eggs laid by the silkworm, *Bull. Seric. Exp. Stn.* **18**(2): 165-171.
- Ganga, G. and Chetty, S.J. (1997). An Introduction to Sericulture, 2nd Edition, Oxford and IBH Publishing Co. Ltd., New Delhi.
- Gatehouse, A. M. R.; Nortan, E.; Davison, G. M.; Babbe, S. M.; Newell, C. A. and Gatehouse, J. A. (1999). Digestive proteolytic activity in leaves of tomato moth, *Lacanobia oleracea*; effects of plant protease inhibitors *in vitro* and *in vivo*. *J. Insect Physiol.* **45**: 545–558.
- Goto, M., Kaneda, M., Imai, T., Murata, S., Fujioka, K., Fujita, E. and Hamamura, Y. (1965). Annual Report of Takeda Research Centre, **24**: 55.
- Govindan, R., Narayanaswamy, T.K. and Devaiah, M.C. (1992). Economic traits and growth indices of eri silkworm, *Samia Cynthia ricini* Boisd., as influenced by substitution of castor with other food plants. *Sericologia* **32** (2):259-263.
- Green, P.W,C, Stevenson P.C., Simmonds, M.S.J. and Sharma, H.C. (2003). Phenolic compounds on the pod surface of pigeonpea, *Cajanus cajan*, mediate feeding behaviour of *Helicoverpa armigera* larvae. *J. Chem. Ecol.* **29**(4):811-821.
- Gülriz Bayçu; Ataa Said; Ahmed Farag and Khaled Rashed (2008). Protein patterns and chemical constituents of *Ailanthus altissima* (Miller) Swingle and *Ailanthus excelsa* Roxb. *IUFS J Biol* **2008**, *67*(1):81-88
- Hamamura Y., Hayashiya K., Naito K., Matsuura K., and Nishida J. (1962). Food selection by silkworm larvae, *Nature*, **194**: 754.
- Hamamura, Y. (2001). Feeding Mechanism and Artificial Diet of Silkworms, In: *Silkworm Rearing on Artificial diet*. (Translated from Japanese), Ed. Yasuji Hamamura, Oxford & IBH Publication, Pp. 1-33.

- 
- Hawksworth, D.L. and Kalin-Arroyo, M.T. (1995). Magnitude and Distribution of Biodiversity. In. V. H. Heywood (ed.). *Global Biodiversity Assessment*, pp. 107-191. Cambrige University Press, Cambrige.
- Haynes. K. F.; Zhao, J.Z. and Latif, A. (1991). Identification of floral compounds from *Abelia grandiflora* that stimulate upwind flight in cabbage looper moths. *J. Chem. Ecol.* **17**: 637-646.
- Hazarika, L.K. and Bhuyan,M. ( 2006). Problems and prospects of non-mulberry germplasm conservation with special reference to the wild silk moths. *Non-mulberry silkworm and host plants: strategies for maintenance*, pp.125-136
- Hazarika, L.K.; Bordoloi, S. and Kataky, A. (1994). Effects of host plants on haemocyte populations and blood volumes of *Antheraea assama* Westwood (Lepidoptera : Saturniidae), *Sericologia*. **34**(2): 301-306.
- Hazarika, L.K.; Saikia, C.N.; Kataky, A.; Bordoloi, S. and Hazarika, J. (1998). Evaluation of physico chemical characteristics of silk fibres of *Antheraea assama* reared on different host plants, *Bioresource Technology*, **64**: 67-70.
- Hazarika, R.L.; Sen, P.; Bhattacharya, S.; Deka, P.C. and Baruah, J.N. (1995) Determination of quality of *Machilus bombycina* for rearing *Antheraea assama* Westwood. *Indian J. Seric.*, **34** (1): 74-75.
- Hegi, G. (1975). in *Illustrierte Flora yon Mitteleuropa*, Vol. 5, pp. 80-87. Verlag Paul und Parey, Berlin.
- Hirao, T. and Ishikawa, S. (1964). Studies on olfactory sensation in the larvae of the silkworm, *Bombyx mori* (I) Chemotaxis of hatched larvae by olfactometer method. *J. Seric. Sci. Jpn.* **33**: 277-285 (in Japanese with English summary).
- Horie, Y. (1978). Quantitative requirements of nutrients for growth of the silkworm, *Bombyx mori* L. *JARQ*. **12**(4): 211-217.

---

House, H.L. (1974). Insect Nutrition, In: *The Physiology of Insecta*. Vol. V (2<sup>nd</sup> Edn.), Rockstein (Ed.), Acad Press. New York, pp. 1-53.

Hughes, J.B.; Daily, G.C. and Ehrlich, P.R. (1997). Population diversity: its extension and extinction. *Science*, 278:689-692.

Ito, T. (1960). Effects of sugars on feeding of larvae of the silkworm, *Bombyx mori*, *J. Insect Physiol.* **5**: 95-107.

Ito, T. (1961 a). Nutrition of the silkworm, *Bombyx mori*, III. Requirements for sterols and their effects on the feeding, *Sanshi Shikenjo Hokoku*, **17**: 91-117. (Original not seen).

Ito, T. (1961 b). Nutrition of the silkworm, *Bombyx mori*, IV. Effects of ascorbic acid, *Sanshi Shikenjo Hokoku*, **17**: 119-36. (Original not seen).

Ito, T. (1961 c). Sterol requirements of the silkworm, *Bombyx mori*, *Nature*, **191**: 882-83.

Ito, T. and Arai, N. (1963). Food values of mulberry for the silkworm, *Bombyx mori* L. determined by means of artificial diets. I. Relationship between kinds of mulberry leaves and larval growth. *Bull. Seric. Exp. Stn.*, Tokyo, **18**(4): 226-229.

Ito, T., and Tanaka, M. (1959). *Per os* administration of the nutrients to the silkworm larvae, I. Effect of the administration of glucose solution, *Sanshi Shikenjo Hokoku*. **15**: 353-64. (Original not seen).

Jat, H. S.; Singh, Ranjay K. and Mann, J. S. (2011). *Ardu* (*Ailanthus* sp) in arid ecosystem: A compatible species for combating with drought and securing livelihood security for resource poor people, *Indian Journal of Traditional Knowledge*, **10**(1): 102-113

Jayaramaiah, M. and Sannappa, B. 1998. Correlation coefficients between foliar constituents of castor genotypes and economic parameters of the eri

- 
- silkworm, *Samiacynthia ricini* Boisduval (Lepidoptera: Saturniidae), *Proc. 3<sup>rd</sup> Intern. Conf. wild silk moths*, Orissa, pp.52.
- Jolly M. S.; Sen, S.K.; Sonowalkar, T.N. and Prasad, G.K. (1979). Non-mulberry silk, Sericulture Manual-4, FAO, Rome.
- Joshi B.C.; Pandey A.; Chaurasia L.; Pal M.; Sharma R.P. and Khare A. (2003b). Antifungal activity of the stem bark of *Ailanthus excelsa*. *Fitoterapia*, 74: 689-691.
- Joshi B.C.; Pandey A.; Sharma R.P. and Khare A. (2003a). Quassinooids from *Ailanthus excelsa*. *Phytochemistry*, 62: 579-584.
- Joshi, K. L. and Mishra, S. D. (1979). Studies on live weight of larvae, cocoons and pupae and silk contents of eri silkworm, *Philosomia ricini* Hull. (Lepidoptera: Saturniidae) reared on interchanged two host plants. *Trans. Indust. & UCDS*, 4(2):97-100
- Joshi, K.L. (1986). Utilization of minerals by larvae of the eri silk moth, *Philosamia ricini* Hutt. (Lepidoptera: Saturniidae), *Indian J. Seric.* 15 (2): 54-57
- Joshi, K.L. (1992). Evaluation of diets for larvae of the eri silkworm *Samia Cynthia ricini* (Boisd) (Lepidoptera: Saturniidae), *Indian J. Seric.* 31 91): 49-51.
- Kakade, M.L. ; Rackie, J.J.; Mcghee, J.E. and Puski, G. (1974). *Cereal Chem.* 51, 376
- Kaleemurrahman, M and Gowri, G. (1982). Foliar constituents of the food plants of eri silkworm (*Philosamia cynthia ricini*). *Proc. Indian natn. Sci. Acad.*, B 48(3):349-353
- Kataky A. and Hazarika L.K. (1997). Fats and fatty acids in the developmental stages of *Antheraea assama* and its host plants, *Sericologia*. 37(3): 511-515.
- Katayama, M.; Holl, C.T.; Chew, N.C.; Hiroto, T.; Kiribuchi, T. and Fanachasi, S. (1974). *Agric. Biol. Chem.*, 38, 1961.

---

Kirtikar K.R. and Basu B.D. (1933). Indian medicinal plants. International Books Distributor, Dehradun, pp 503-507

Kirtikar, K.R. and B.D. Basu (1995). Indian Medicinal Plants. Vol. 1, International Book Distributors, Dehradun, India, pp: 371-372.

Kohli, R.K.; Jolly, M.S.V. and Khan, A.M. (1969). Foliar constituents of the food plants of tasar silkworm *Antheraea mylitta* D., *Indian For.* **95**(9): 614-617.

Koopowitz, H. and Kaye,H. (1990). *Plant Extinction –a global crisis*. Christopher Helm, London.

Krishnaswami, S.; Noamani, M.K.P. and Asan, M. (1970). Studies on the quality of mulberry leaves and silkworm cocoon crop production. I. Quality difference due to varieties. *Indian J. Seric.* **9**: 1-10.

Kumar ,D.; Bhujbal S.S.; Deoda, R.S. and Mudgade, S.C. (2010). Bronchodilator activity of aqueous extract of stem bark of *Ailanthus excelsa* Roxb. Phcog Res, 2:102-6

Kumar, Dinesh; Bhat, Z.A.; Singh, P.; Shah, M.Y. and Bhujbal,S.S. (2010). *Ailanthus excelsa* Roxb. is Really a Plant of Heaven, International Journal of Pharmacology 6 (5): 535-550

Kumar, R and Elangovan, V (2010). Rearing Performance of Eri Silkworm, *Philosamia ricini* in Monsoon Season of Uttar Pradesh, *Asian J. Exp. Biol. Sci.*, 1 (2 ): 303 – 310.

Kumari, S. Saritha; Narayanswamy, K.C. and Gowda, Manjunath (2009). Amylase Activity in Eri Silkworm, Samia Cynthia ricini Boisduval as influenced by Host Plants. *Indian J. Ecol.* (2009)36(1) : 71-74.

Kundu, Pijush and Laskar, Subrata (2010). A brief resume on the genus Ailanthus: chemical and pharmacological aspects. *Phtochem Rev.* 9: 379-412

---

---

Lavhale, M.S. and Mishra,S.H. (2007). Nutritional and therapeutic potential of *Ailanthus excelsa*: A review. *Pharmacognosy Rev.*, 1: 105-113.

Lewis, N.G. and Yamamoto, E. (1990). Lignin: occurrence, biogenesis and biodegradation. *Ann. Rev. Plant Physiol. Plant Mol. Biol.* 41: 455-496.

Li, Hui-lin (1993). "Simaroubaceae". in Editorial Committee of the Flora of Taiwan. *Flora of Taiwan, Volume 3: Hamamelidaceae-Umbelliferae* (2nd ed.)  
<http://tai2.ntu.edu.tw/udth/bin/fot1.exe/browse>

Li, R. and Sanu, O. (1984). The relationship between quality of mulberry leaves and some economic characters during the later larval stage. *Canye Kexue*, 10(4): 197-201.

Lin S., Binder B.F. and Hart E.R. (1998). Insect feeding stimulants from the leaf surface of *Populus*. *J. Chem. Ecol.*, 24(11): 1781-1790.

Loizzo, M.R.; Said A.; Tundis, R.; Rashed K.; Statti, G.A.; Hufner A. and Menichini, F. (2007) Inhibition of Angiotensin Converting Enzyme (ACE) by Flavonoids isolated from *Ailanthus excelsa* (Roxb)(Simaroubaceae). *Phytotherapy Research*, 21: 32-36

Lopez, J. D.; Shaler, Jr.. T. N.; Beerwinkl, K. R. and Lindgren, P.D. (2000). Feeding attractant and stimulant for adult control of noctuid and/or other lepidopteran species, US Patent 6,074,634, Issued June 2000, United States Patent Office, Washington, D.C.

Loschiavo, S. R. (1965). The chemosensory influence of some extracts of Brewer's yeast and cereal products on the feeding behavior of the confused flour beetles, *Tribolium confusum* (Coleoptera: Tenebrionidae). *Ann. Entomol. Soc. Am.* 58: 576-588.

Malick, C.P. and Singh, M.B. (1980). In: *Plant Enzymology and Histo Enzymology*, Kalyani Publishers, New Delhi, Pp. 286.

---

Mallette, M.F.; Althous, P.M. and Clagetta, C.O. (1960). *Biochemistry of Plants and Products*. Wiley Eastern Pvt. Ltd., New Delhi.

Maxwell-Lefroy, H. and Ghosh, C.C. (1912). Eri Silk. Memoirs of the Department of Agriculture in India, Entomological Series, 4 (1):130

May, R.M.; Lawton, J.H. and Stork, N.E. (1995). Assessing extinction rates. In: *Extinction Rates* (eds. Lawton, J.H. and May, R.M.). Oxford University Press, London. pp.1-24

Medical Economics Company (1998). PDR for herbal medicines, *Ailanthus altissima*. Montvale, New Jersey, p 618.

Merenjungla and Kakati, L.N. (2013). Seasonality of nutrient contents of different leaf types of four host plants of *Samia ricini* Donovan in Nagaland, *National Journal of Life Science*, Vol. 10(2): 105-112

Michael Eskin, M.A.; Hochn, E. and Frenkel, C. (1978). A simple and rapid quantitative method for total phenol, *J. Agric. Food Chem.*, 26: 973.

Ministry of Environment and Forests, Government of India F.C. Division, New Delhi (2004). Office Order under Forest (Conservation) Act, 1980 No. F.No. 2-1/2003-FC (Pt II) dated 7<sup>th</sup> June 2004.

Myers, N.; Mittermeier, R.A.; Mittermeier, C.G.; Fonseca, G.A.B. and kent, J.( 2000). Biodiversity hot spots for conservation priorities, *Nature*, 403:853-858

Nadkarni K.M. (1976). Indian material medica. Bombay Popular Prakashan, Bombay, p 56.

Nag, A. and Matai, S. (1994) *Ailanthus excels* Roxb. (Simaroubaceae), a Promising Source of Leaf Protein. *Journal of Agriculture and Food Chemistry*, 42: 1115-1117.

---

---

Narayana Prakash, M.; Periswamy, K. and Radhakrishnan, S. (1985). Effect of dietary water content on food utilization and silk production in *Bombyx mori* L. (Lepidoptera: Bombycidae), *Indian J. Seric.*, 24(1): 12-17.

Narayanaswamy, K. C.; Kumari, K. S.; Manjunath Gowda and Amarnatha, N. (2006): Barkesseru Ailanthus excelsa Roxb., an ideal substitute for castor to rear eri silkworm, *Samia cynthia ricini* Boisduval. *Environment and Ecology* 24(4): 720-725

Neilson J.K., Larson L.M. and Soresen H. (1979). Host plant selection of horseradish flea beetle *Phyllotreta armoraciae* (Coleoptera: Chrysomelidae): Identification of two flavonols glycosides stimulating feeding in combination with glucosinolates, *Entomol. Exp. Appl.*, 26:40-48.

Neog, K. (2011). Study on biochemical stimulants in host plants for developing an effective indoor rearing technique of muga silkworm (*Antheraea assamensis* Helfer) (Lepidoptera: Saturniidae). Ph.D. Thesis, Gauhati University, Guwahati.

Neog, K; Das, A.; Unni, B.G.; Ahmed, G.U. and Rajan, R.K. (2011a). Studies on Secondary metabolites of Som (*Persea bombycina* Kost), a primary host plant of Muga silkworm (*Antheraea assamensis* Helfer), *Int. J. PharmTech Res.* 3(3): 1441-1447

Neog, K; Unni B,G and Ahmed ,G.( 2011b). Studies on the influence of host plants and effect of chemical stimulants on the feeding behavior in the muga silkworm, *Antheraea assamensis*. *Journal of Insect Science* 11:133

Nicholson R and Hammerschmidt R. (1992). Phenolic compounds and their role in disease resistance. *Ann. Rev. Phytopathol.* 30: 369-389.

Nooteboom HP. 1962. Simaroubaceae. In: Van Steenis CGGJ, ed. Flora Malesiana ser. 1 vol. 6. Groningen: Wolters-Noordhoff Publishing, 193–226.

- 
- Ogura M.; Cordell G.A.; Kinghorn A.D. and Farnsworth N.R. (1977). Potential anticancer agents VI. Constituents of *Ailanthus excelsa* (Simaroubaceae). *Lloydia*, 40:579-584
- Pandey, R.K. (1995). Do leaf tannins affect non-mulberry silkworm, *Indian Silk* 34 (8): 21-23.
- Parpiev, B.A. (1968). Water metabolism in silkworm fed with a different mulberry strain changing diet. *Shelk.* 39: 15-17.
- Pathak, A.K. (1988). Studies on nutrition, growth and cocoon characters of eri silkworm *Philosamia ricini* Hutt. fed on different varieties of leaves. M.Sc, Thesis, Assam Agri. University. Jorhat (India), pp.18-65
- Paul D.C.; Subba Rao G. and Deb, D.C. (1992). Impact of dietary moisture on nutritional indices and growth of *Bombyx mori* and concomitant larval duration. *J. Insect Physiol.* 38(3): 229-235.
- Peigler, R. (1992). *Ibid*, 32(3):191
- Peter P. Feret (1985). Ailanthus: variation, cultivation, and Frustration. *Journal of Arboriculture* 11 (1 2): 361
- Phelan, P. L.; Roelofs, C. J.; Youngman, R. R. and Baker, T. C. (1991). Characterization of chemicals mediating ovipositional host-plant finding by *Amyelois transitella* females. *J. Chem. Ecol.* 17: 599-613.
- Phukan, J.C.D.; Singh, K.C.; Neog, K. and Chakravorty, R. (2006). *Ailanthus grandis* prain (Simaroubaceae- Quassia family) An alternate food plants of eri silkworm *Samia ricini* (DONOVAN). In: Proceeding of National workshop on Eri food plants. 2005 Guwahati, 102-110.
- Prasong,S.; Yaowalak, S.and Wilaiwan, S. (2009). Characteristics of Silk Fiber with and without Sericin Component: A Comparison between *Bombyx mori* and *Philosamia ricini* Silks. *Pakistan Journal of Biological Sciences*, 12: 872-876.

---

Rai, S.; Agarwal, K.K. and Babu, C.R. (2006). Influence of host plant (*Terminalia arjuna*) defences on the evolution of feeding behaviour in the tasar silkworm, *Current Science*. **91**(1): 68-72.

Rajanandh M.G. and Kavitha J. (2010). Quantitative Estimation of  $\beta$ -Sitosterol, Total Phenolic and Flavonoid Compounds in the Leaves of *Moringa oleifera*, *Intern, J. Pharm Tech. Res.*, **2**(2): 1409-1414.

Ravikumar (1988). Western ghats as a bivoltine region: Prospects, challenges and strategies for its development. *Indian Silk* **26** (11): 39-54.

Roark, B. and J. E. Quisenberry (1977). Environmental and genetic components of stomatal behaviour in two genotypes of upland cotton. *Plant Physiol.* **59**: 354-356.

Sadasivam, S., Manickam, A. (2005). Biochemical Methods. Newage International Pvt. Limited, New Delhi.

Sarkar A. A.; Haque M. R.; Rab M. A. and Absar, N. (1997) Studies on crude protein and amino acid contents of mulberry (*Morus alba* L.) leaves in relation to cocoon production of silkworm *Bombyx mori*. *Sericologia*. **30**(1): 17-22.

Sarkar, D.C. (1980). *Sericulture in India*. 1<sup>st</sup> edn. (Bombay: Central Silk Board).

Sarmah ,M.C.; Rahman, S.A.S.; and Chakravorty, R. (2006). Biodiversity of Eri Host Plants in North east India-Its conservation and constraints. Non-Mulberry Silkworm and Host Plants Germplasm. pp.11-17

Sarmah, M. C., Debaraj, Y., Singh, B. K., Das, P. K. and Suryanarayana, N. (2002). Status paper on eri host plant Germplasm, Proceedings: National workshop on Sericultural Germplasm Management and Utilization, Hosur, Tamil Nadu, 6-7th February 202 P: 59-61

Sarmah, M.C., Neog, K., Das, A, Phukan, J.C.D., (2013). Impact of soil fertility and leaf nutrients status on cocoon production of Muga silkworm, *Antheraea*

---

---

---

*assamensis* (Helper) in potential muga growing areas of Assam, India,  
*Int.J.Curr.Microbiol.App.Sci* 2(9): 25-38

Sarmah, M.C.; Chutia, M.; Neog, K.; Das, R.; Rajkhowa, G. and Gogoi, S.N. (2011). Evaluation of promising castor genotype in terms of agronomical and yield attributing traits, biochemical properties and rearing performance of eri silkworm, *Samia ricini* (Donovan). *Industrial Crops and Products*, 34 (2011):1439– 1446

Sarmah, M.C.; Sarkar, B. N.; Ahmed, S. A. and Deuri, J. (2013a). Ericulture: A comprehensive profile, Directorate of Sericulture, BTC, Kokrajhar.

Sarmah, R.K. (1992). Studies on the xternal morphology and commparative rearing performance of eri silkworm, *Philosamia ricini* Hutt. (Lepidoptera: Saturniidae)on different host plants, M.Sc. Thesis, AAU, Jorhat.

Schanderl, S.H. (1970). In: *Methods in Food Analysis*, Academic Press, New York, Pp. 709.

Schoonhooven, L. M. (1982). Biological aspects of antifeedants. *Entomol. Exp. Appl.* 31: 57-69.

Scriber, J.M. and Slansky, F. Jr. (1981). The Nutritional Ecology of Immature Insects, *Annual Review of Entomology*, 26: 183-211

Sharma, P.V. (1996). Dravya Guna-Vijnana, ayurvedic series-III, vegetable drugs. Chaukhamba Bharati Academy, Varanasi, pp 466-468.

Shaw, C. (1998). Evaluation of *Ailanthus* species in relation to nutrition, growth and cocoon characters of eri silkworm, *Philosomia ricini* Hutt., M. Sc. Thesis, AAU, Jorhat.

Sherman, M.M.; Borris, R.P.; Ogura M.; Cordell,G.A. and Farnsworth, N.R. (1980) 3S,24S,25 Trihydroxytirucall-7-ene from *Ailanthus excelsa*. *Phytochemistry*, 19:1499-1501.

---

Shrimali M.; Jain D.C.; Darokar, M.P. and Sharma, R.P. (2001). Antibacterial activity of *Ailanthus excelsa* (Roxb). *Phytotherapy Research*, 15: 165-166.

Simmonds M.S.J. (2001). Importance of flavonoids in insect-plant interactions: feeding and oviposition. *Phytochemistry*, 56: 245-252.

Singh, N. I. & Goswami, D. 2012. Food plant varietal effect on the rearing and grainage of Muga Silkworm, *Antheraea assamensis* Helfer, 1837. *Munis Entomology & Zoology*, 7(2): 1023-1027

Sinha, A.K. and Jolly, M.S. (1971). Foliar constituents of the food plants of tasar silkworm, *Antheraea mylitta* D., *Indian For.* 97(5): 262-263.

Sinha, A.K.; Choudhury, S.K.; Brahmacharai, B.N. and Sengupta, K. (1986). Foliar constituents of the food plants of temperate tasar silkworm, *Antheraea proylei*, *Indian J. Seric.*, 25(1): 42-43.

Sinha, U. S. P.; Neog, K.; Rajkhowa, G. and Chakravorty, R. (2003). Variations in biochemical constituents in relation to maturity of leaves among seven mulberry varieties. *Sericologia*. 43(2): 251-257.

Sinha, U. S. P.; Sinha, A.K.; Srivastava, P.P. and Brahmacharai, B.N. (1992). Studies on the variations in chemical constituents in relation to maturity of leaves in three primary food plants of tropical tasar silkworm, *Antheraea mylitta* D., *Indian J. Seric.*, 31(1): 83-86.

Slansky F., Jr. and J.M. Scriber (1985). Food consumption and utilization. pp. 87–163 in G.A. Kerkut and L.I. Gilbert, eds. *Comprehensive Insect Physiology, Biochemistry, and Pharmacology*. Vol. 4. Pergamon, Oxford.

Sticher, L., Mauch-Mani, B., and Metraux, J.P. (1997). *Ann. Rev. Phytopathol.* 35, 235.

Takeuchi, Y. (1960). Ability of silkworm (*Bombyx mori* L.) to recover from malnutrition. *The Silk News*. 5(8): 6-7.

- 
- Tamaki, Y.; Loschiavo, S. R. and McGinnis, A. J. (1971). Triglycerides in wheat germ as chemical stimuli eliciting aggregation of the confused flour beetle, *Tribolium confusum* (Coleoptera: Tenebrionidae). *J. Agr. Food Chem.* **19**: 285-288.
- Tanaka, Y. (1964). *Sericology* (Bombay: Central Silk Board)
- Tangamani, R. and Vivekanandan, M. (1984). Physiological studies and leaf nutrient analysis in the evaluation of best mulberry variety, *Sericologia*, **24**: 317-324.
- Thorsteinson, A.J. (1960). Host selection in phytophagous insect, *Ann. Rev. Entomon.*, **5**: 193-218.
- Torii, I and Morii K. (1948). Studies on the feeding habits of silkworms, *Reports of the Silk Science Research Institute*, (SANSHIKA): **2**: 3.
- Unni B.G., Kakoty A.C., Khanikar D., Bhattacharya P.R., Pathak M.G., Pillai K.P.R., Choudhury A., Saikia A. and Ghosh A.C. (1996). Lipid and fatty acid composition of muga silkworm, *Antheraea assama* host plants in relation to silkworm growth, *J. Lipid Mediators and Cell Signalling.*, **13**: 295.
- Urzua, A. (2002). Monoterpenes and sesquiterpenes in the headspace volatiles from intact plants of *Pseudognaphalium vira vira*, *p. heterotrichium*, *p. cheiranthifolium* and *p. robustum*: their insect repellent function, *Bol. Soc. Chil. Quím.* **47**: 099-104.
- Van Soest, Peter J. 1982. Nutritional ecology of the ruminant: ruminant metabolism, nutritional strategies, the cellulolytic fermentation, and the chemistry of forages and plant fibers. Corvallis, Oregon: O & B Books.
- Vance, C.P., Anderson, J.O., and Sherwood, R.T. (1976). *Plant Physiol.* **57**, 920.
- Vasuki K. and Basavanna H. M. (1969). Variety difference in the content of total and soluble minerals of mulberry leaves. *Silkworm Information Bulletin.* **I**: 31-35.

---

Verk, J. S., Ravinder Kaur and Parwinder Kaur (2009). Performance of eri silkworm, *Samia ricini* Boisduval in different seasons of Punjab. Indian J. Seric., 2009, 48(1):78-80.

Verschaffelt, E. (1910). The cause determining the selection of food in some herbivorous insects. *Proc. Royal Acad. Amsterdam.* **13:** 536–542.

Waldbauer, G.P. (1968). The consumption and utilization of food by insects. Advances in Insect Physiology 5:229–288.

Watanabe, T. (1958). Substances in mulberry leaves which attract silkworm larvae (*Bombyx mori*). *Nature.* **182:** 325-26.

Wheeler, E.L. and Ferrel, R.E. (1971). A method for phytic acid determination in wheat and wheat fractions. *Cereal Chem.* **48:** 312-315.

Williams, N.H. (1981). The Biology of Orchids and Euglossine Bees. In *J. Arditti ed. Orchid Biology, Reviews and Perspectives II.* Cornell Univ. Press, Ithaca New York, pp. 119-171.

Wilson, E.O. (1992). *The Diversity of Life.* Norton, New York.

Y. Chen (1933). Silviculture of Chinese trees. 20 + 656 + 4 + 6 pp. Nanking (In Chinese.)

Yates, I. E., Bacon, C. W., and Hinton, D. M. (1997). Plant Dis. 81 (7), 723

Yoshida, T (1955). Recent advances in silkworm nutrition, *Bull. Sericult. Expt. Sta.* **14(7):** 351-426

Zheng, Hao; Wu, Yun; Ding, Jianqing; Binion, Denise; Fu, Weidong; Reardon, Richard (2004). "Ailanthus altissima" (PDF). *Invasive Plants of Asian Origin Established in the United States and Their Natural Enemies, Volume 1.* USDA Forest Service.