

## BIBLIOGRAPHY

1. Achimu, P. and Schloesse, E. (1992). Effect of neem seed extract *A.indica* against downy mildew of grapevine. *Int. Symp. Crop. Prot.*, 2<sup>nd</sup> & 4<sup>th</sup> May, Gent (Belgium), **44**(2).
2. Adhikari, R.S. and Tiwari, A. (1991). Some experimental studies of phylloplane and litter fungi of *Quercus semicarrufolia*. *J.Ind.Bot. Soc.*, **70** (129-134).
3. Adhikari, R. S. (2002). Phylloplane mycoflora of three grasses of *Pithoragarh*. *Acta Bot. Indica*.**26**: 285-291.
4. Ahmed, S.A., Dutta, L.C., and Sarmah,M.C. (2012). Bioefficacy of some insecticides against leaf eating caterpillar *Cricula trifenestrata* Helfer (Lepidoptera : Saturniidae) infesting Som *P.bombycina* Kost. Plantation. *Academic journal of Entomology*. **5**(2): 94-98
5. Ahmed, S.A. and Rajan, R.K. (2011). Exploration of Vanya silk biodiversity in NER of India : A sustainable livelihood and poverty alleviation. *Int. Con. On Management, Economics & Social Sc. (ICMESS, 2011)*
6. Ahmed, N. and Grainge, M. (1982). Some promising plants for pest control under small scale farming operation. *Working paper resource system*. Institute East West Centre Honolulu, Howai, U.S.A.: 342-346
7. Amadioha, A.C. (1998). Fungitoxic activity of extracts of *Azadirachta indica* and *Xylopi aethiopica* on *Cottetotrichum lindemuthianum* in Cowpea. *Journal of Herbs, Spices and Medicinal Plants*, **6**: 33-40.
8. Amadioha, A.C. (2003). Fungitoxic effects of extracts of *Azadirachta indica* against *Cochliobolus miyabeanus* causing Brown Spot Disease of Rice. *Acta. Phytopath. Pflanz.* (Taylor & Francis), **35**: 37-42.
9. Anonymous (1962). The wealth of India. Council of Scientific and Industrial Research, New Delhi. India. **6**: 429-439.
10. Anonymous (1981). Muga Silk Industry, Directorate of Sericulture & weaving, Govt. of Assam, Guwahati.
11. Anonymous (2010). Climate change poses threat to silk production in Assam, Newspaper report in *Assam Times*.
12. Ansari, A.R. and Prakash, D. (1986). Rhizosphere and rhizoplane mycoflora of barley infected with *Ustilago hordie*. *Ind. Phytopath.* **39**:380-384

13. Anuratha, C.S. and Gnanamanickam, S.S. (1987). *Pseudomonas fluorescences* suppresses development of bacterial blight symptoms. *IRRN*. **12**(1): 17
14. AOAC(1975). Official methods of Association of Agricultural chemists. Hartwitz W. (ed.), Washongton, 12<sup>th</sup> edition.
15. Al-Abed , A.S., Qasem,J.R. and Abu-Blam,H.A. (1993). Antifungal effects of some common wild plants species on certain plant pathogenic fungi. *Dirasat* (Pure and Applied Science), **20**: 149- 158.
16. Atlas, R.M., and Parks, L.C. (1997). Handbook of microbial media. 2<sup>nd</sup> ed. C press; Boca Raton.
17. Babu, L.J., Gour, H.N. and Pankaj, S. (2008). Efficacy of phytoextracts and bioagents against *Colletotrichum gloeosporioides* causing banana fruit rot. *J. Mycol. Pl. Pathol.* **38**: 635-637.
18. Barah, A., Goswami, M.C. and Samson, MV. (1989). Consumption & utilization of food in different instars of muga silkworm *A.assama* WW. *Proc. Indian Acad Sci. (Anim Sci)*, **98**(2):99
19. Barah, A., Singh, P.K. and Sengupta, A.K. (1992). Studies on preservation on muga cocoons at low temperature. *Ann. Report* (1991-92), RMRS, CSB, Boko, Assam.10-11.
20. Barnard, C., Padgitt, M., and Uri, N. D. (1997). Pesticide use and its measurement *Intern. Pest Control.* **39**:161-164
21. Barnes, G. (1971). Inhibition of *Erysiphe polygoni* on clover leaf surface by saprophytic spores in “Ecology of leaf surface microorganisms” edited by Preece and Dickinson (A.P.London)
22. Barua, A. (1991). Studies on certain aspects of the activity of the soil and seed mycoflora of Sal forest in Kamrup district. *Ind. Phytopath.*, **60**:386-390.
23. Baruah, A. and Bara, K. (1995). Seasonal variation and distribution of microfungial population in Sal tree forest soil in Kamrup distric Assam. *Annual Technical Session o f Assam Science Society*, Guwahati. Assam. **16**.31.
24. Barnett, H.L. (1960). Illustrated Genera of Imperfect Fungi . 2<sup>nd</sup> Ed, Published by Burgess Publishing Co.

25. Barsagade, N.B. and Wagh, G.N. (2002). Comparative screening of leaf extracts of common plants and weeds for their antibacterial and antifungal activities. *AJOBR*, **3**: 227-232
26. Basavanna, H.M. and Bose, P.C. (1989). Characteristics of soil of mulberry farm of Central Sericultural Research & Training Institute, Mysore. *Indian J. Seric.* **28**(1):1-10
27. Bennet, S.S.R. (1987). Name changes in flowering plants of India and adjacent regions. Triseas publishers, Dehradun.
28. Bernstein, T.B. and Feinberg, S.M. (1942). Airborne fungus spores. A five year survey of daily mold spore content of Chicago air. *J.Allergy*, **13**:231
29. Bhattacharjee, Kathakalee., Sharma, G.C. and Kalita, S. (2012). Aeromycological study in a bus terminus in guwahati city, Assam. *AJEBS*. **3**(2): 311-319
30. Bhattacharyya, PN. and Jha, DK. (2011). Seasonal and depthwise variation in microfungus population numbers in Nameri forest soil, Assam, North east India. *Mycosphere*, **2**(4): 297-305
31. Bharali, N. (1969). Grey blight disease of som. *Plant Sci. Cult.*, **35**: 573-574
32. Bharali, N. (1985). Facts on sericulture in Assam. Directorate of sericulture, Govt. of Assam.
33. Bhowmik, T.P. (1974). Fungicidal control of *Alternaria* leaf blight of wheat. *Indian Phytopath.* **27**: 162-167
34. Bhuyan,P.M., Sandilya,S.P., and Gogoi,D.K., (2013). Phyllosphere microflora of muga silkworm host plant *P.bombycina* Kost. (Som) in leaves in Jorhat district of Assam India. *IRJBS*, **2** (12): 60-65
35. Bilgramij, S.S. and Jamaluddin, Rizwi, M.A. (1979). Fungi of India. 1. Published by Today and Tomorrow's printers and publication. New Delhi. 121-145.
36. Biswas, I. And Ray, N. (2006). Effect of environmental factors on seed (egg) production of muga silk worm (*A.assama* Ww) in terrain region of West Bengal. *Indian Journal of Entomology*. **68**(1):39-43
37. Biswas, S., and Sing, N.P., (2005). Fungicidal management of foliar disease of groundnut in Tripura. *Ind. Phytopath.***84**:500-502
38. Black, C.A., (1965). Method of Soil analysis.1., Am.Soc. Agron., Madison Wisconsin, USA

39. Blakeman, J. P. and Brodie, I.D.S. (1976). Inhibition of pathogens by epiphytic bacteria on aerial plant surface in "Microbiology of Aerial plant surface" edited by Preece & Dickinson.
40. Bongale,U.D. (1993). Fertility evaluation and fertiliser recommendations for mulberry garden soils. *Indian Silk*. **31**(9): 5-8
41. Bongale,U.D. and Chaluvachari, K. (1995). Evaluation of eight mulberry varieties by leaf biochemical and biosafety moulting studies. *Sericologia*. **35**(1): 83-94
42. Bongale,U.D. and Lingaiah. (1998). Macro and micronutrient status of mulberry garden soils in a bivoltine seed area. *Indian J. Seric.*, **37**(1):73-75
43. Bongale, U.D. and Siddalingaswamy,K. (1996). Physicochemical properties of soil from major sericultural areas of Kolar and Bangalore districts , Karnataka, India. *Indian J. Seric.* **35**(2): 152-154
44. Bora, A., Borah, BR., Sengupta, AK. (1990). Studies on preservation of muga cocoons of higher altitude. *Ann. Report RMRS, CSB, Boko, Assam.*15-16
45. Bora, L.C., Gangapodhya, S. and Chand, J.N., (2004). Phylloplane colonizer of three mung bean cultivars in relation to growing season and plant age. *Ind. Phytopath.***64**:203-208
46. Bora,A., Shukanand,SN., and Samson,MV. (1988). Variation in cocoon characters of *A.assama* WW during different seasons. *Sericologia*, **28** :215
47. Bordoloi, D.N., and Baruah,H.K.(1964). Studies on distribution of certain fungi in tea plantation soil and air. *J. Univ.G.U. Sci.* **17**(1): 155-170
48. Borgohain, A., Das,R., and Chutia,M., (2014). Fungal diversity in phylloplane of castor plant (*Ricinus cummunis* L.): the primary food plant of Eri silkworm. *Scholarly Journal of Agricultural Science*. **4** (2): 82-86
49. Borthakur, S.K. (2003). Ethnobiological wisdom behind the traditional muga silk industry in Assam. *Ind J. of Trad. Knowledge*,**2**(1): 230-235
50. Borthakur, B., Mitra, J.B., and Gogoi, P. (2001). Rhizosphere mycoflora of *Aquilqria agattocha* Roxb. *Proceeding of Seminar on Scope and dimension of Agar plantation in N. E. region.*34-37
51. Bray, R.H., and Kurtz, L.T. (1945). Determination of total organic and available forms of Phosphorus in soils. *Soil Science* ,**59**: 39-45

52. Buragohain, R. (1994). The Lost Trails (Ban-ok pup-lik Mioung-Tai, Dhemaji). **1**.
53. Bussaman, P., Namsena, P., Rattanasena, P. and Chandrapatya, A. (2012). Effect of crude leaf extracts on *Colletotrichum gloeosporioides* (Penz.) Sacc.. *Psyche*.1-6
54. Chakravorty,R., Barah,A., Neog,K., Das,K. and Kakati,P.K. (2006). Variability in muga silkworm *Antheraea assamensis* Helfer. *Indian Silk*. 16-17
55. Chakravorty, R., Das, R., Neog,K., Das,K. and Sahu,M.(2007). A diagnostic manual for diseases and pests of muga silkworm and their host plants. Central Muga and Eri Research & Training Inst., Lahdoigarh, Jorhat.
56. Chakraborty, S., Perrott,R., Charchar,MJ., Fernandes,CD., and Kelumu,S. (1997). Biodiversity, epidemiology, and virulence of *Colletotrichum gloeosporioides*. II. Genetic and pathogenic diversity in isolates of *Colletotrichum* from eight species of *Stylosanthes*. *Tropical Grasslands*. **31**: 393-401.
57. Chand JN, Kondal MR, Aggarwal RK (1968). Epidemiology and control of bitter rot of apple caused by *Gloesporium fructigenum* Berk. *Ind Phytopathol*. **21**: 257-263.
58. Chandwani, G.H., Blakrishnanan, M.S., and Padmanabhan, S.Y. (1963). *Helminthosporium* disease of rice. V. A study of the spore population of *H.oryzae* over rice fields. *J. of Indian Botanical Society*. **42**:1-4.
59. Chattopadhyay, C. (1999). Yield loss attribute to *Alternaria* blight of sunflower (*Helianthus annuus L*) in India and some potentially effective control measures. *Inter. J. Pest Manage*. **45**(1): 15-21
60. Chaudhuri, M., Sing , SS., Das, B., Dhar, NJ., Basumutary, B., and Goswami, D. (1999). Climatic variability in nine locations of NE India and their effect on cocoon productivity of muga silkworm (*A.assama* westwood). *Sericologia*. **39** (4):577-591.
61. Chiang, H.C. (1985). Insects and their environment, Fundamentals of applied Entomology, Macmillan Publishing Company, New York. 128-161.
62. Choudhuri, S.N. (1981). Muga silk industry, Directorate of sericulture and weaving, Govt. of Assam, Gauhati, Assam, India. 1-33.
63. Choudhury, SN. (1970). Muga Silk Industry, Directorate of Sericulture, Govt. of Assam, Dispur, Guwahati, 23.
64. Choudhury, S. N. (2005). Host plant species of non-mulberry (vanya) silkworm. *Proc. Work. Str. Non. Mul. Germ. Main.*, CMER&TI, Jorhat Assam. 50-54.

65. Choudhury, M., Singh, S.S., Das, B., Dhar, N.J., Basumatary, B., Goswami, D., Das, K., Barua, A., Sahu, M., Kakoty, L.N., Mandal, T., and Chatterjee, S.N. (1999). Climatic variability in nine locations of north east India and their effect on cocoon productivity of Muga silkworm (*Antheraea assama* Westwood). *Sericologia*, **39**(4): 577-591
66. Chowdhuri, S.N. (1980). Reeling and weaving of muga silk. *Indian Silk*. **19**(5):15
67. Chowdhury, MAM, Auda, MK., and Iseam ASMT (2000). Phytodiversity of *Dipterocarpus tubinatus* Gaertn.F. (Garjan) undergrowths at Dulahazara garjan forest. Cos' B Bazar, Bangladesh. *Indian Forester*. **126**:674-684
68. Chowdappa, P., Reddy, G.S., Kumar, A., Rao, B.M., and Rawal, R.D. (2009). Morphological and molecular characterization of *Colletotrichum* species causing anthracnose of grape in India. *The Asian Aus. J. Pl. Sc. Biotech.*, **3**: 71-77.
69. Cordier, T. Robic., Capdevielle, X., Desprez Loustau, M.L., and Vachor, C. (2012). Spatial variability of phyllosphere fungal assemblages: genetic distance predominates over geographic distance in a European beech stand (*Fagus sylvatica*), *Fungal Ecology*, **5**(5):509-520
70. Cox, L.A., and Hall, A. M. (1978). Phylloplane fungi of *Quercus robur*. *Proceedings of Annals and Applied Biology*. **89**:119-122.
71. Cumminghumb, D.D. (1873). Microscopic examinations of air. Government printer, Calcutta, India, 58.
72. Curtin D., Campbell C.A., Zenter R.P., and Lapond G. (1994). Long term management and clay dispersibility in two haploborolls in Saskatchewan, *Soil Sci. Soc. Am. J.* **58**(3):962-967
73. Dalal, L.P. (2012). Incidence and diversity of soil mycoflora of Wardha (M.S) area. *Int. J. of Life Sc. and Pharma Res.*, **4**(4): 15-20
74. Dalton, E.T. (1872). Descriptive ethnology of Bengal (Supdt. Govt. Printing, Calcutta)
75. Das, R., and Benchamin, K.V. (2000). Diseases of muga and eri food plants : Incidence pattern, Intensity & control measures. *Proceedings of National seminar on sericulture, R & D in muga & eri*, CMER&TI, Jorhat Assam, 34-35
76. Das, R., Das K. and Rajan, R.K. (2014). Effect of plant extract on foliar blight disease of soalu caused by *Colletotrichum gloesporioides* Penz. **3**(2):76-80
77. Das, K. K., and Dubey, L.N. (2001). Studies on rhizosphere mycoflora of agar plant in Assam. *Proceeding of Seminar on Scope and dimension of Agar plantation in N. E. region*. Pp. 6-12.

78. Das, K., and Das, R. (2003) Controlling of Grey blight disease of som plant. *Ind. J. Seric.* **40**:139-143.
79. Das, K., and Das, R., Baruah, P. and Bara, K. (2005). Studies on leaf surface mycoflora in soalu plant in different seasons. *Proceedings of 49th Annual Technical Session, Assam Science Society*, **5**: 198-205.
80. Das, R., Chutia, M., Das, K., and Jha, D.K. (2010). Factors affecting sporulation of *P. disseminata* causing grey blight disease of *Persea bombycina* Kost., the primary food plant of muga silkworm. *Crop Protect.*, **29**: 963-968
81. Das, R. and Jha, D.K. (2008). Evaluation of some chemicals against *P. disseminata* causing grey blight of som. *Sericologia*. **48**(1): 113-117
82. Das, R., Das, K. and K.V. (2003). Diseases of muga food plant som. *Indian Silk*. **41**:23-24
83. Deb, P.R., Deb, M. and Dutta, B.K. (1999). A preliminary report on phyllosphere mycoflora of tea and the soil mycoflora of an experimental tea plantation area of Cachar. *Indian Phytopath.* **52**(2): 193-195
84. Debnath, S. and Baruah, P.K. (2008). Seasonal variation of air mycoflora over tea plantation in Jorhat district, Assam. *Ind. J. Aerobiol.* **21**(2): 79-85
85. Deshmukh, A.J., Mehta, B.P., Sabalpara, A.N., Patil, V.A. (2012). Screening of Indian bean (*Lablab purpureus* L.) varieties/ germplasms against anthracnose (*Colletotrichum gloeosporioides* Penz. and Sacc.) under field conditions. *J. Biopest.*, **5**: 50-52.
86. Devi, O.J., and Chetry, G.K.N. (2013). Evaluation of antifungal properties of certain plants against *Drechslera oryzae* causing brown leaf spot of rice in Manipur valley. *IJSRP*. **3**(5): 1-3
87. Devanath, M; Ramesh, A; Ramanujam, K., and Ramabadrhan, R. (1994). Evaluation of fungicides for the control of leaf blight of Guava, caused by *Alternaria solani*. Diseases of Horticulture plants. Kalyani Publishers, New Delhi- 2. 511-513.
88. Dickinson, C.H. and Preece, T.F. (1976). "Microbiology of Aerial plant surface" A.P. London.
89. Dongmo, J.C., and Oyeyiola, G.P., (2009). *Fusaria* in the root zone of tomato (*Lycopersicon esculentum*). *Nigerian Journal of Microbiology*. **23**(1):1922-1932

90. Dubey,R.C., and Dwivedi,R.S. (1991). Fungitoxic properties of some plant extracts against vegetative growth and sclerotial viability of *Macrophomina phaseolina*. *Indian Phytopath.*, **44**(3): 411-413
91. Dubish,P. K.; Srivastava,S.; Sing,L., and Bala, S., (1989). Rhizosphere mycoflora of virus infected and healthy plants of *Lagenaria vulgaris* and *Abelmoschus esculantus*. *Ad. Pl. Sci.* **2** : 191-196.
92. Dutta,B.K.,(1981). Studies on some fungi isolated from the rhizosphere of tomato plants and the consequent prospect for the control of *Verticillium* wilt. *Plant and Soil.***63**(2): 209-216
93. Dutta,L.C., Kalita,M.N., and Sarkar,C.R., (1997). Foliar constituents of the food plants of muga silkworm *A.assama* WW. *Indian J. Sericulture.* **36**(1):85-86
94. Dutta, S; Dutta,B.K., and Nath,P.K. (2010).Comparative study of air, phyllosphere and soil mycoflora of the tea plantation area of Cachar district, Assam. *Assam University Journal of Science & Technology: Biological and Environmental Sciences*, **5**(1): 89-94
95. Dutta,S.K., Roychoudhury,S., pandit,D., and Bajpai,A.K., (2010). Rot diseases of muga host plant som (*P.bombycina*) in Assam, India. *The journal of plant protect. Sci.* **2**(2): 73-76
96. Erpelding,J.E., and Prom,L.K., (2004). *Plant Pathology Journal.***3**:65-71
97. Fokunang,C.N., Ikotun,T., Dixon,A.G.O., Akem,C.N., Tembe,E.A., and Nukenine,E.N., (2000). Efficacy of antimicrobial plant crude extracts on the growth of *Colletotrichum gloeosporioides* f. Sp. manihotis. *Pakistan Journal of Biological Sc.*, **3**(6):928-932.
98. Fonseca,T.C., De Almeida, J.E., okamoto, F.,and da Cunha,E.A. (1993). Nutritive value of some mulberry clones (*Morus alba* L.). *Sericologia.***33**(4): 631-633
99. Forsyth,F.R., and Peterson,B.,(1960). Control of leaf and stem rust of wheat by Zineb and inorganic nickel slats. *Pl. Dis. Reprtr.* **44**:208-211
100. Funk,DT.; Neely ,D., and Bey,CF.,(1981). *Silvae Genetica.* **30**: 4-5.
101. Gadre,U.A., Sharma,I.P. ,and Mandokhot,A.M., (1990). Studies on population threshold of some pathogenic fungi on mango in relation to weather parameters in Kongkan region. *A paper presented at seminar on strategies for increasing mango production at Konkan Krishi Vidyapith, Dapoli, India on 22<sup>nd</sup> and 23<sup>rd</sup> March.*



102. Ganapaty,T.,and Narayanaswamy ,P.(1994). Screening plant products effective against Tikka and Rust diseases of Groundnut. *Crop Diseases, Innovative Techniques and Management*. Kalyani publishers, New Delhi-110002. 348-353
103. Gangawane ,L.I. and Kulkani ,L. (1985). Rhizosphere mycoflora of groundnut grown in sewage and sludge treated Soils. *Ind. Phythopath.* **38**:756-758.
104. Gangawar,S. K.; Maji,M.D.; Kumar,R .,and Sarathchandra, B., (2002). Evaluation of fresh plant extracts for control of mulberry powdery mildew. *Ind. J. Seric.* **39** :76-79.
105. Gangwar,S.K., Quadri,M.H., Maji,M.D., Kumar,P., and Saratchandra,B. (2000). Evaluation of fresh plant extracts for the control of mulberry powdery mildew. *Indian J.of Seric.* **39**(1): 76-78.
106. Ganie,S.A., Pant ,V.R., Ghani,M.Y., Lone,A.H., Anjum,H., and Razvi,M. (2013). Invitro evaluation of plant extracts against *A.brassicae* (Berk.) Sacc. causing leaf spot of mustard & *Fusarium oxysporum* f. Sp. lycopersici causing wilt of tomato. *Academic Journal.* **8**(37):1808-1811.
107. Garg,A. P.,and Sharma,P.D. (1987). Phylloplane mycoflora of non-infected and rust infected barley and triticale. *Acta Bot. Indica.* **8** : 57-60
108. Ghosh, SK. And Chakraborty, N. (2012). *Invitro* biological control of *Colletotrichum gloeosporioides*, causal organism of Anthracnose of Sarpagandha (*Roulvolfia serpentina*). *Agric. Biol. J. N. Am.* **3**(8):306-310
109. Giorgio,C. di.; Krempff,A.; Guiraud,H., and Dumentl,G.,(2006) Atmospheric pollution by air borne microorganisms in the city Marseilles. *Atmospheric Environment* **48**:198-206.
110. Gilman, C. Joseph. (1995 reprint). A manual of Soil fungi. Printwell, Jaipur.(India).
111. Gogoi,A.K., Singh,G.P., Sahu,A.K., and Giridhar,K. (2013). Screening of foliar diseases tolerant varieties of Som, *Persea bombycina*- a primary food plant of muga silkworm, *Antheraea assamensis* Helfer. *Souvenir cum compendium of 1<sup>st</sup> Bodoland National Seminar cum Workshop on Sericulture industry & its role in the socio-economic upliftment in rural society.*
112. Gogoi,B. (1996). *Muga culture*- The heritage of *Tai-Ahoms* in Assam, *Souvenir*, (Purbanchal Tai Sahitya Sabha, Assam)
113. Gogoi, B. And Goswami, B.C. (1998). Studies on rearing performance of muga silkworm (*Antheraea assama* Ww) on Dighloti (*Litsea salicifolia* Roxb.) *Sericologia.* **38**(3): 511-514

114. Goswami,C., and Bhattacharya,Manisha., (2013). Contribution of sericulture to women's income in Assam- A case study in Goalpara district of Assam, India. *Int. J. of Sci. & Res. Publications*; **3**(3).ISSN 2250-3153
115. Govindaih,Gunashekar.V. and Thiagarajan.(2002). Field evaluation of fungicides against powdery mildew in mulberry plant *Indian J. of Seric.***33** (2): 160-162
116. Govindaiah. Phillip T., Sengupta,K., Giridhar,K. and Suijyanarayana,N.(1990). Incidence of leaf disease in relation to epidermal and cutical thickness in some indigenous varieties of mulberry. *Sericol.* **30** : 257-260.
117. Goncalves,PDS., Furtado,EL., Batagila,OC., Ortolani,AA., May,A. and Bellerri,G.O., (1999): *Genetics Molecular Biology.* **22**: 583-589.
118. Gunashekar,V. and Govindaiah (1999). Monitoring of mulberry diseases. *Ind. Silk .* 50-52.
119. Gunashekar,V., Govindaih and Himantharaj,M.T., (1995). Effect of fungicides in controlling mulberry leaf rust caused by *Cerotelium fici*. *Indian J. of Seric.* **34**(1): 60-62.
120. Gupta,VK., Pandey,A., Kumar,P., Pandey,BK., and Gaur,RK. (2010). Genetic characterization of mango anthracnose pathogen *Colletotrichum gloeosporioides* Penz. by random amplified polymorphic DNA analysis. *Af. J. Biotech.* **9**: 4009-4013.
121. Hackel,E., Zechmeister,B.S., Bodrossy,L., and Sessitsch,A.(2004). Comparison of diversities and compositions of bacterial populations inhibiting natural forest soils. *Applied & Environmental Microbiology.* **70**:5057-5065
122. Harde,A.L. and Suryawanshi,A.P. (2014). Evaluation of some antifungal plant extracts against *A.brassiccae*, inciting *Alternaria* blight of mustard. *Trends in biosciences.* **7**(11): 1007-1011
123. Hardison,J.R. (1963). Commercial control of *Puccinia striiformis* and other rusts in seed crops of *Poa pratensis* by nickel fungicides. *Phytopathology.* **53**:209-216
124. Hamilton, E.D.(1968). A comparison of the pollen and fungus spore contest of air in two localities and concentration of the respiratory allergy. Ph.D. Thesis, University press. London.
125. Harlapur,S.I., Kulkami,M.S., Wali,M.C., and Srikant,Kulkmil.(2007). Evaluation of plant extracts, bio agents and fungicides against *Exserohilium turcicum* (Pass) Leonard and Suggs. causing *Turcicum* leaf blight of maize. *Karnataka J. of Agri. Sci.* **20**(3): 541-544.

126. Harsh,N.S.K., Nath,V., Tiwari,C.K., and Rehill,P.S.,(1987). Studies on a new foliar disease of *Diospyros melanoxylon*. Roxb. *Van Vigyan*. **25**:16-20.
127. Hazarika,R., Hazarika,L.K., Kakaty,A., Kakaty,J.C.S., Hazarika,A.K., and Deka,P.C. (1996). Association of morphological and biochemical characters of *Machilus bombycina* with the feeding behavior of *Antheraea assama*. *Sericologia*. **36**:511-518
128. Hazarika, R.L., Sen, P., Bhattacharya, S., Deka, P.C., and Baruah, J.N. (1995). Determination of quality of *Machilus bombycina* for rearing *A.assama* Westwood. *Indian J. Seric*. **34**(1): 74-75
129. Hegde,YR. and Hegde,RK. (1986). Studies an anthracnose of arecanut (*Areca catechu* L.) caused by *Colletotrichum gloeosporioides* (Penz.) Penz. and Sacc. *Pl. Pathol. Newslett*. **4**: 24-27
130. Hooker,J.D.(1885). The flora of British India. William clowes and Sons Ltd., London; **5**:155-180.
131. Hossain,I., and Schloesser,E.,(1993) Control of *Bipolaris sorokiniana* on wheat with neem extracts. *Bangaladesh J Microbiol*. **10**: 39-42.
132. Hubballia,M., Nakkeerana,S., Raguchandera,T., Ananda,T. and Renukadevi,P. (2011). Physiological characterisation of *Colletotrichum gloeosporioides*, the incitant of anthracnose disease of noni in India. *Arch. Phytopathol. Pl. Prot*. **44**:1105-1114.
133. Huq,M.I.(2011). Studies on the epidemiology of leaf rot and leaf spot diseases of Betel vine (*Piper Betle* L.). *Bangladesh J. Sci. Ind. Res*. **46**(4): 512-522.
134. Hyd,H. A. and William,W.A. (1953). The incidence of *Cladosporium herbarum* in the out door air at Cardiff. *Tram. Brit. Myco. Soc*. **41**: 165-167.
135. Iqbal, M.C.M., Meiyalaghan,S., Bwijesekara,K., and Abeyratne,K.P. (2001). Antifungal activity from water extracts of some common weeds. *Pakistan Journal of Biological Sc*. **4** (7): 843-845
136. Islam,M.R., Hossazin ,M. K., Bahar,M. H. and Ali,M.R. (2004). Identification of the causal agent of leaf spot of Betelnut and *in vitro* evaluation of fungicides and plant extracts against it. *Pakistan J. of Biol. Sci*.**7**:1758-1761.
137. Ivarson, K.C. and Mack, AR. (1972). Root surface mycoflora of soybean in relation to soil temperature and moisture in a field environment. *Can. J. Soil. Sci*. **52**:199-208

138. Iyenger ,M. N. S.(1995). Effects of feeding mulberry leaves sprayed with Morestan on the survival and growth of silkworms. *Bombyx mori. Technical Session 6, Second Forestry Conference.*
139. Jackson, M.L. (1958). Soil chemical analysis. Prentice Hall of India Private Limited, New Delhi, 1978
140. Jalander,V., And Mamatha,M., (2015). Rhizosphere mycoflora of some leguminous crop plants. *Int. J. Pure App. Biosci.* **3**(3): 262-266
141. Jaya Prakash, P., Rao, S.B.V. Sanjeev, and Rao, J.V. Krishna (2004). Scope for muga culture in the agency are as of Visakapatnam district of Andhra Pradesh. *Proceedings of National workshop, CSB, Dehradun.* 300-303.
142. Jespers,A. B. K., and Ward,M.A. de. (1993). Natural products in plant protection. *Netherland J. Pl. path.* **99** :109
143. Jeyalakshmi,C. and Seetharaman,K. (1998). Biological control of fruit rot and dieback of chilli with plant products and antagonistic microorganisms. *Pl. Dis. Res.***13**: 46-48
144. Jolly ,M.S., Chowdhury, S.N. and Sen,S.K.(1975). Non-mulberry sericulture in India. Central Silk Board, Bombay. 25.
145. Joshi,M.C., Singha, P.S., Kumar,R. S., Beck and Singh, B.R.P. (2004). Occurrence of foliar diseases in vanya silkworm food plants and their management. *International Journal of Pest Management.* **51**: 15-21.
146. Jorgensen,L.N. and Olsen,L.V.(2007). Control of tan spot (*Drechslera tritici-repentis*) using different fungicides. *Crop Prot.* **26** : 45-50.
147. Kadam, J., Kadam, J.J., Sudhir Navathe and Agale, R.C. (2014). Epidemiology of leaf blight disease of turmeric caused by *Colletotrichum gloeosporioides*. (Penz. And Sacc.). *Scholarly J. Agric. Sci.* **4**:470-475
148. Kagale,S.; Marimuthu,T.; Nanda Kumar, R., and Samiyappan,R. (2007). Antimicrobial activities of leaf extract of *Datura metela* against *Rhizoctonia solani*. *Physiol. Mol Pathol.* **65** : 91-100
149. Kagty ,L.C., (1964). Influence of soil types on the distribution of fungi in the rhizosphere of wheat. *J. Ass. Sci. Soc .I*: 11-19.
150. Kahle,P., Leinweber,P., and Menning,P., (1992). On the influence of soil organic matter on physical properties of soils. *Agric. Biol. Res.* **45**(1):18-27
151. Kakati, L., and Hazarika,B.N.,(1997). Significant variation in lipid content in muga food plant som. *Ind. J. Seric.* **44** : 269-272.

152. Kamanna, B.C. (1996). Epidemiology & control of Anthracnose diseases of coffee incited by *Colletotrichum gloeosporioides* (Penz.) Penz & Sacc. PhD Thesis, UAS, Bangalore, 185.
153. Kanappa, BC. (1998). Epidemiology and control of anthracnose of diseases of coffee incited by *Colletotrichum gloeosporioides* (Penz.) Penz. and Sacc. University of Agricultural Sciences, Bangalore, India
154. Kanjilal,U.N., De,P.C., and Das, R.N., (1992). Nyctaginaceae to cycadaceae. Flora of Assam. **4**, 66-67
155. Kasprzyk,I., Rzepowska,B., and Wasylow,M. (2004). Fungal spores in the atmosphere of Rzeszow (South East Poland). *Ann. Agricul. Envi. Medicine*. **11** : 285-289.
156. Kasprzyk,I. and Worek ,M. (2007). Airborne fungal spores in urban and rural environments in Poland. *Aerobiologia*. **22** : 169-176.
157. Kaushal,Aarti. and Sing, Ranjana. (2013). Diversity and seasonal variation of soil fungi isolated from surrounding area of upper lake, Bhopal, Madhya Pradesh. *Adv. In Life Sc. & Tech*.**8**:1-3
158. Keith,LM., Velasquez,ME. and Zee, FT. (2006). Identification and characterization of *Pestalotiopsis* spp. causing scab disease of guava, *Psidium guajava* in Hawaii. *Plant Dis*. 90:16–23
159. Khaleqizaman,K., Hossain,M.L., and Hossain,M.M. (2001). Effect of fungicides & potash in controlling grey leaf spot of coconut. *Bangaladesh journal of trading and development*. **11**: 151-156
160. Khanna,KK. and Chandra,S .(1976). Studies on storage diseases of fruits and vegetables. Control of guava fruit rot caused by *Pestalotia psidii* and *Colletotrichum gloeosporioides* with Aretan. *Curr. Sc*. **45**: 70.
161. Khanikar, D. and Unni,B.G. (2006). Food plant biochemistry and its role in growth and development of muga silkworm, *A.assama* WW. *Muga silkworm Biochemistry, biotechnology Molecular biology*.88-101.
162. Khatri, R.K. (2003). Effect of seasonal variation on muga silkworm (*A.assama* Ww) rearing and grainage under different conditions of Doon valley. Ph.D. Thesis, HNB Garhwal University, Sri Nagar, Garhwal.
163. Khatri, R.K., Singh, N., Paliwal, D., Lal, Babu., Siddiqui, A.A. and Sharma, A.K.(2004). Correlation studies on environmental factors and rearing and grainage behaviour of *Antheraea assama* Westwood under Doon valley conditions. *National workshop on potential and strategies for sustainable development of Vanya silks in the Himalayan states, Dehradun, Uttarakhand*.160-166

164. Kim,M., Sing,D., Lai-Hoe,A., GoR., Rahim,R.A., Ainuddin,A.N., Chun,J. and Adams ,J.M. (2012). Distinctive phyllosphere bacterial communities in tropical trees. *Microb Ecol.* **63**(3): 674-681.
165. Kofalvi, S.A., Gao, J. G. and Nassuth (2007). Biochemical investigation of wheat leaves infected by mosaic virus. *Physiol. & Mol. Pl. Pathol.* **58** : 376-389.
166. Konger,G. and Barua, H. K. (1970). The incidence of air borne spores in the potato plantation of upper Shillong. *J. Gua. Univ.* **9** : 81-84.
167. Krishnakumar,A.K., Dutta,A.K. and Potty,S.N. (1990). Moisture retention characteristics of soils under Hevea in India. *Indian J. Rubber Res.* **3**(1): 9-21
168. Krishnaswami S., Noamani M.K.R. and Ahsan,M. (1970). Studies on quality of mulberry leaves and silkworm cocoon crop production I. quality differences due to varieties. *Ind. J. of sericulture.* **9**: 1-10.
169. Kuberan,T., Balamuruga, A., Vidhyapallavi, R., Nepolean,P. and Jayanthi,R. (2012). In Vitro Evaluation Certain Plant Extracts against *Glomerella cingulata* Causing Brown Blight Disease of Tea. *World J. Agricul. Sc.* **8**: 464- 467.
170. Kulkarni, Pragya. (2011). Aeromycological profile of the public parks of Bhilai Township, Chhattisgarh, India. *Indian Journal of Science and Technology.* **4**(5): 558-560
171. Kumar, A., Sharma,JN., and Garg,R.C. (2012a). Prevalence, symptomatology, pathogenicity and nutritional requirement of *Colletotrichum gloeosporioides* causing premature fruit drop in Kinnow. *Ind. Phytopathol.* **65**: 366-372.
172. Kumar,S., Thind,TS., and Mohan,C. (1995). Morphogenic and pathogenic variations in *Gloeosporium ampelophagum*. *Ind. Phytopathol.* **48**: 331-334.
173. Kumer,N.U. and Gupta,J.S. (1976). Phyllosphere mycoflora of three potato varieties in relation with different weather factors. *Ind. Phytopath.* **42** : 116-120
174. Kundalkar,S.K., Joshi,M.S., and Pawar,D.R. (1991). Control of leaf blight of coconut caused by *Pestalotia palmarum* CKE. *Indian coconut journal.***22**:18-19.
175. Kurian,PS., Sivakumar,G., Josephraj Kumar,A., Backiyarani,S., Murugan,M. and Shiva,K.N. (2008) .Management of anthracnose disease (*Colletotrichum gloeosporioides* (Penz) Penz & Sac.) of black pepper (*Piper nigrum* L.) in the high ranges of Idukki District, Kerala. *J. Spices Aromatic Crops.* **17**: 21-23.
176. Lal,R.(1977). Physical properties and moisture retention characteristics of some Nigerian soils. *Int. Inst. trophi. Agri.* **8**(1),1350

177. Last, F.T.(1955). The spore content of air within and above mildew infested cereal crops. *Trans. Br. Mycol. Appl.* **44**: 289-294
178. Last, F.T. and Deighton, F.C. (1965). The non parasitic microflora on the surface of living leaves. *Transactions of the British Mycological Society.* **48**: 83-99
179. Langenberg, W. J., Sulton, J.C. and Gillespie (1977). Relation of weather variability and periodicities of air borne spores of *Alternaria dauci*. *Ind. Phytopath.* **67**:879-883
180. Leben, C. (1965). Epiphytic microorganisms in relationship to plant diseases. *Annl. Rev. Phytopathology.* **3** : 209-230
181. Leveau, J. (2009). Life on leaves, *Nature.* **461**: 741
182. Majumder, S. K., Bose, P.C., Banejee, N.D. and Parthesh, Kumar P.M. (2003). Ascorbic acid, an index of fungal infection of mulberry leaves. *Sericologia.* **43**:119-122.
183. Manjunath. (2009). Morphological and Molecular characterization of *Alternaria alternata* and *Colletotrichum gloeosporioides* incitants of leaf blight and anthracnose diseases of Noni and their Management, Tamil Nadu Agricultural University, Coimbatore, India
184. Manoharachary, C., Vankataswalu, K. and Ramarao, P. (1977). Studies on rhizosphere and non-rhizosphere soil of Kharif plants. *GEOBIOS.* **4**: 67-68.
185. Mdee, L.K., Masoko, P. and Eloff, J.N. (2009). The activity of extracts of seven common invasive plant species on fungal phytopathogens. *South African J. of Bot.* doi: 10.1016/j.sajb.2009.02.003
186. Mitra, S.R., Chaudhury, A. and Choudhury, N. (1994). Production of antifungal compound by higher plants. A review of recent researches. *J. Physiol. Biochem.* **11**:53-90.
187. Mohanan, C.R. And Kaveriappa, K.M. (1982). Occurrence and distribution of Cacao diseases in Southern India. *Proceedings of the 8<sup>th</sup> International Cocoa Conference* . 445-450
188. Mohanan, R.C., Kaveriappa, K.M. and Nambiar, K.K.N. (1991). Variability in growth of *Colletotrichum gloeosporioides* isolates pathogenic on cacao in response to fungicides, antibiotics and detergent. *J. Plantation Crops.* **18**: 229-232.
189. Mondali, N.K., Mojumdar, A., Chatterje, S.K., Banerjee, A., Datta, J.K., and Gupta, S. (2009). Antifungal activities and chemical characterization of neem leaf extracts on the growth of some selected fungal species in *in vitro* culture medium. *J. Appl. Sci. Environ. Manage.* **13**(1): 49-53

190. Morris,C., and Kinkel,L. (2002). Fifty years of phyllosphere microbiology: significant contributions to research in related fields. *Phyllosphere microbiology*, Lindro, S., E. Hecht-poinar and V.Elliott, (Eds.) APS press, St. Paul, MN, USA. 365-375.
191. Mukhopadhyas,D. and Nandi, B. (1977). Rhizosphere mycoflora of healthy and diseased jute plants. *Curr. Sci.* **43**: 590-593
192. Muthukrishnan, J. And Pandian, T.J. (1987). Relation between feeding and egg production in some insects. *Proc. Indian Aca. Sci. (Anim. Sci.)*.**96**(3):171-179
193. Nagaraja,T.G. (1990). Rhizosphere mycoflora of *Strychnos nux-vomica*. *Ind. Bot. Reprtr.* **9**:32-33
194. Nene, Y.L. and Thapliyal, P.N. (1993). Evaluation of fungicides. Fungicides in Plant Disease control. Oxford and IBH Publishing company, New Delhi.531.
195. Neog, K., Chakravorty, R. and Golgi, SN. (2007). Evaluation of som (*Persea bombycina*) germplasm accessions through bioassay for rearing performances and biochemical analysis of leaves. *Proceedings of Sericulture challenges in the 21<sup>st</sup> century (Serichal 2007) and 3<sup>rd</sup> BACSA meeting.* 81-87.
196. Old, KM., See, Lee ,Su., Sharma,JK., and Yuan, Zi Qing. (2000). A manual of disease of tropical acacias in Australia, South East Asia and India. Center for International Forestry Research, Jakarta, Indonesia and CSIRO Forestry and Forest Products Canberra, Australia.
197. Oyeyiola,G.P. (2009). Rhizosphere mycoflora of Okro (*Hibiscus esculentus*). *Res. Journal Of Soil Biol.* **1**(1): 31-36
198. Oyelia,G.P.(2010). Rhizosphere bacterial flora of *A.hybridus*. *Res. J. of Microbiology*.**5**(2):137-143
199. Padule,D. N., and Hande,Y.K.(1980). Laboratory evaluation of some fungicides against *Pestalotia psidii* the cause of guava canker. *Pesticides.* **14**:37-38.
200. Pal, GK., and Kumar, B. (2013). Antifungal activity of some common weed extracts against wilt causing fungi, *F.oxysporum*. *Int. J. of curr. Discoveries and innovations.* **2**(1):62-67.
201. Pal, GK., Kumar, B., and Shahi, S.K. (2013). Antifungal activity of some common weed extracts against seed borne phytopathogenic fungi *Alternaria sp.* *Int. J. of Univ. Pharmacy and Life Sc.* **3**(2): 6-14.



202. Pandey, A. (2011). Variability studies and molecular characterization of *Colletotrichum gloeosporioides* causing anthracnose of mango (Ph.D. thesis). A.P.S. University, Rewa, M.P. India.
203. Pandey,A., Sukla,A.N., and Chandra, S. (2006). *Pestalotiopsis* stem canker of *Jatropha curcas*. *Indian Forester*. **132**: 763-766.
204. Pandey,R.K., Singha,U. N. and Goel,P.K. (1993). Constituents of young and old leaves of primary food plants of oak tasar silkworm *Antheraeaaproyle*. *Ind. J. Seric*. **30**:70-71.
205. Pandey,R.K., and Dwivedi,R.S.(2000). Seasonal incidence of Phylloplane mycoflora of guava with reference to the pathogens. *Acta Bot. Indica*. **22**: 185-189.
206. Pandey,A., Kamle,M., Chauhan,UK. and Pandey,BK .(2009). Evaluation of plant extracts against *Colletotrichum gloeosporioides* an incited of mango anthracnose disease. *Pl. Arch*. **9**: 947-949.
207. Patel, DS. (2009). Chemical management of fruit spot of pomegranate caused by *Colletotrichum gloeosporioides* Penz. and Sacc. *Ind. Phytopathol*. **62**: 252- 253.
208. Patel,RV. and Joshi, K.R. (2002). Efficacy of different fungicides against *Colletotrichum gloeosporioides* Penz. and Sacc. causing leaf spot of turmeric. *J. Mycol. Pl. Pathol*. **32**: 413-414.
209. Patel,SI., Patel,JI. and Vadodaria,JR. (2007). Efficacy of fungicides against leaf and fruit spots of pomegranate caused by *Colletotrichum gloeosporioides*. *J. Mycol. Pl. Pathol*. **2**: 379-340.
210. Patil, C. R., and Chandra, A. K. (1980). Variation in the number of phyllospheric nitrogen fixtures and other microplane in relation to the age of the host plants. *Curr. Sci*. **49**:655-657.
211. Patil, C. R., and Thite, A.N. (1987). Rhizosphere mycoflora of four hybride varieties of *Sorgham vulgare*. *Ind. J. Mycol. Pl. Path*. **17**:50-56
212. Philip, T. and Govindaiah (1995). *Pestalotiopsis disseminata* (Turn) Steyaert- A new pathogen on mulberry. *Indian J. of Seric*. **34**(2): 159-160.
213. Phukan,R., and Chowdhury,S.N. (2006). Traditional knowledge and practices involved in muga culture of Assam. *Indian journal of Traditional knowledge*. **5**(4): 450-453.

214. Powelson,R.L., and Shanner,G.E. (1966). An effective chemical seed treatment for the systematic control of seedling infection by strips rust (*Puccinia striiformis*). *Pl. Dis. Reprtr.* **50**:806-807
215. Potphode,P.D., and Fugro,P.A. (2012). Epidemiology and management anthracnose of bitter gourd (*Momordia charntia*) incited by *Colletotrichum lagenarium* (Pass.) Ell. and Halst. *Indian Phytopathology.***65**: (3-S) -18
216. Prasad, G.K. and S.S. Sinha (1982). Induction of diapauses in muga silkworm. *Annual report, CMER&TI, Jorhat, Assam.*
217. Prashanth,A., Arun, RS., Naik ,M.K., Patil, M.B. and Rajesh, S.P. (2008). Evaluation of fungicides, bioagents, and botanicals against pomegranate anthracnose. *Ind. J. Pl. Prot.* **36**: 283-287.
218. Prince, L., and Prabhakaran, P. (2012). Studies on the soil mycoflora from the sugarcane field in Thanjavur district, Tamilnadu. *J. Microbiol.Biotech. Res.* **2**(1): 63-69
219. Pugalmaran,M. and Vittal, BPR. (2000). Fungal diversity in the indoor and outdoor environments and dust grain storage godowns. *Ind J Aerobiol.* **12**:24-29
220. Qasem, Abu-Blam, H.A. (1996). Fungicidalactivity of some common weed extracts against different plant pathogenic fungi. *Journal of Phytopathology.* **144**:157-161
221. Quadri,M.D., Gangwar, S.K., and Prathees, K. (2004). Efficacy of fungicides in controlling mulberry foliar diseases. *Indian J. of Seric.* **34**(2):18-21.
222. Rahman,A., Tanti,B., Sharma, G.C. and Kalita, J.(2012). Genetic diversity of *P.bombycina* from Goalpara district of Assam, India. *Adv. In BioSc. & Biotechnology,* **3**: 20-24
223. Ramesh,C.R., Chakraborty,S. and Sukanya,D.H. (2004). Role of *Stylosanthes fruticosa* in diversity of *Colletotrichum gloeosporioides* in India. Hibh- yielding antracnose- resitant Stylosanthes for agricultural systems. Australian Center for International Agricultural Reasearch (ACIAR) Canberra.
224. Ramesh,C.; Kabalgari,B.R.,and Shahny,L., (2003). Seasonal variation and diversity of air borne fimgi o f *Lavender* plantation in Dharward. Karnataka. *J Mycol. Pl. Pathol.*,**32** : 34-38
225. Ramesh,C., Patil,L.,and Shameembanu,K.(2002). Population dynamic of Microfungi in rhizosphere region of cotton with special reference to seasonal variation in Dharward. *J. Mycol. Pl. Path.* **32** :10-17.

226. Rampersad, SN. (2011). Molecular and phenotypic characterization of *Colletotrichum* species associated with anthracnose disease of papaya in Trinidad. *Plant Dis.* **95**:1244-1254
227. Rangaswami,G., and Bagyaraj, D.J. (1988). Agricultural microbiology, 2nd edition, published by prentice hall of India Pvt. Limited.
228. Rao,KS. and Satyanarayana, A. (1989). Epidemiology of anthracnose of grape (*Vitis vinifera*) caused by *Elsinoe ampelina* around Hyderabad. *Ind. J .Agricul .Sc* .**59**: 655-657.
229. Ray, Manjit Kumar., and Mishra, P.K. (2012). A preliminary Screening and identification of microflora in the outdoor and indoor environment of B.N.College, Dhubri, Assam. *Ann. Plant Physiol.* **26**(1): 35-39
230. Reddy, K.M.,and Ramanujam, C.GK.(1989). An Aerobiological study of Hydrabad (AP). *Asian J. Pl.. Sci.*, **11**:7-21.
231. Reddy, G. Vemananda, Vijayalakshmi, Rao., and Samson, M.V.(1999). Role of embryonic stage during long term preservation of bivoltine silkworm. *Bombyx mori* C. eggs on hatchability. *Abs. In: Breeders Scientist interaction issues related to germplasm maintenance, protection and utilisation, 10<sup>th</sup> Feb, CSGRC, Hosur.*
232. Ruinen, J. (1961). The phyllosphere-an ecologically neglected milliew. *Plant and Soil.* **15**:81-109
233. Saikia, P. and Chowdhury, H.D. (1993). Phylloplane microflora for the control of bacterial leaf blight of rice caused by X.O.PV. *Oryzae. Indian phytopath.* **46**(3):218-223
234. Saikia, S. and Goswami, B.C. (1997). Mejankari: The unexplored Assam Silk. *Indian Silk.* **35**(12): 33-35
235. Sahu, A.K. and Bindroo, B.B.(2007). Wild silk moth biodiversity in the North Eastern Region of India: need for conservation. *Indian Silk.* **46**:16-19
236. Saju KA, Deka TN, Gupta U, Biswas AK. and Sudharshan MR. (2012). *In vitro* evaluation of biocontrol agents, botanicals and fungicides against *Colletotrichum gloeosporioides* infecting large cardamom. *Pl. Dis. Res.***27**: 49-53.
237. Saju,KA., Mech, Smrita., Deka, TN., and Biswas, AK.( 2011). In vitro evaluation of biocontrol agents, botanicals and fungicides against *Pestalotiopsis sp.* Infecting large cardamom (*Amomum subulatum* Roxb.). *Journal of Spices and Aromatic crops.***20**(2):89-92
238. Saksena, HK. and Kumar, K. (1985). Proceeding of Indian National Science Academy. **37**:399-406.

239. Samanta, A., Chatterjee, A.K., Kar, R., and Mandal, B. (2001). Assessment of manganese content in mulberry garden soils of West Bengal. *Indian J. Seric.*, **40**(1):64-70
240. Sanguri, S., Kapil, S., Gopinathan, P., Pandey, F.R. and Bhatnagar, T. (2012). Comparative screening of antibacterial & antifungal activities of some weeds & medicinal plants leaf extracts: an invitro study. *Elixir Appl. Botany*. **47**:8903-8905.
241. Sankhla, B., Sankla, H.C., Dalela, G.G., and Mathur, R.L. (1972). Evaluation of fungicides against blight disease of wheat caused by *Alternaria triticina*. *Indian Phytopath.* **25**:210-214
242. Santhi, D.P., Unnamalai, N. and Gnanamanickam, S.S.(1987). Epiphytic association of *Erwinia herbicola* with rice leaves infected by X.C.PV. *Oryzae* and its interaction with pathogen. *Indian Phytopath.* **40**(3):327-332
243. Saraf, A., Quereshi, S., Sharma, K. and Khan, N.A. (2011). Antimicrobial activity of *Lantana camara* L. *Journal of Experimental Sci.* **2**(10):50-54.
244. Saratchandra, B.(2006). Muga silk industry - Its status and strategies, Muga silkworm Biochemistry Biotechnology Molecular-biology published by RRL, Jorhat, Assam, India.161-167.
245. Sarkar, S.K. and Samaddar, K.R. (1978). Occurrence & interactions between aerial plant surface microorganisms. *Ind. Sci. Cong. Ari. Sect.* **65**: B (Abstract)
246. Sarkar, A.A., M.R. Haque, Rab, M.A. and Absar, N. (1997). Studies on crude protein & amino acid contents of mulberry (*Morus alba* L.) leaves in relation to cocoon production of silkworm *Bombyx mori*. *Sericologia*. **30**(1): 17-22.
247. Sarvamangala, H.S.; Govindaiah and Dutta, R.K.(1993). Evaluation of plant extracts for the control of fungal diseases of mulberry food plant. *Ind. Phytopath.* **46** : 398-401.
248. Saxena, A.K., (2002). Anthracnose of pomegranate-Biology of the pathogen, Epidemiology and disease control. PhD thesis, submitted to Maharshi Dayanand Saraswathi University, Ajmer, Rajasthan, India.
249. Saxena, S.B. (1954) . Ecological factors governing distribution of soil microfungi in some forest soil of Sagar. *J. Ind. Bot. Soc.* **24**:262-298.
250. Sengupta, R.R., Das, K., Devnath, M. and Samson, M.V. (1993). Collection, identification & evaluation of *Machilus bombycina* King. (Laurales: Lauraceae) germplasm, the muga food plant. *Sericologia*. **33**:109-110
251. Sengupta, K., Singh, K.(1974). Some studies on the effect of refrigeration of muga *A.assamensis* cocoons on moth emergence, egg laying and hatching of eggs. *First International Seminar on non-mulberry silks, CTR&TI, Ranchi*.174

252. Sengupta, A.K., Srivastava, A.K., Raja Ram, and Negi, B.K.(1995). Muga- new strides. *Indian Silk*. **35**(3):28-32
253. Seth, M.K. (2000). Food plant of muga silkworm. In: Agarwal, H.O. & Seth, M.K., Eds., *Sericulture in India*, Bishen Singh Mahendra Pal Singh, Dehradun. **4**:887-893.
254. Shahnaz, E., Razdan, V. K., and Raina, P.K. (2007).Survival, dispersal and management of foliar blight pathogens of onion (*Allium cepa*). *J. of Mycol. and Plant Pathol.* **37**(2): 210-214.
255. Shahny, M., and Purwar, A.(2002). Studies on occurrence of aeromycoflora in Allahabad University campus. *Ind. J. Aerobiol.* **16**: 24-29.
256. Shampatkumar, A., Eswara, Reddy NP., Hariprasad Reddy, K., Chowdappa, P., Subhash Reddy, G .(2007). Genetic Diversity and Pathogenic Variability among *Colletotrichum gloeosporioides* Penz Isolates: The Causal Agent of Mango Anthracnose. *ICFAI. J. Biotec.* **2**: 15-27.
257. Sharma, A. K., Gupta, J. S .,and Dixit, R. B.,(1992). A quantitative and qualitative analysis of Phylloplane mycoflora yellow sarson and tamarina in relation to microclimatic factors. *Ind. Phytopath.* **40**:109-113
258. Sharma, A. and Verma ,KS. (2007). In vitro cross pathogenicity and management of *Colletotrichum gloeosporioides* causing anthracnose of Mango. *Ann. Pl. Prot. Sc.* **15**: 186-188.
259. Sharma, B., and Kumar,P.(2014). In vitro antifungal potency of some plant extracts against *F.oxysporum*. *Int. J. of Green Pharmacy.*63-65.
260. Sharma, D. ,and Dutta, B. K.(2002). Observation of aeromycoflora of jute plantation in Silchar area. *Ind. J. Aerobiol.***16**:9-16.
261. Sharma, D. K. and Devi, D. (1997). Seasonal Variation of the Foliar Constituents of the Primary Food Plants of the Muga Silkworm (*Antheraea assama* W/w). *Sericologia.* **37**(2): 251-258.
262. Sharma, G. D.,and Tiwari, B. K. (1988). Leaf surface mycoflora of healthy and diseased leaves (*Phytophthora infestans*) *Solanum khasianum*. *Acta Botanica Indica.* **9**:233-237
263. Sharma, IM., Raj ,H., Kaul ,JL. And Raj, H. (1994). Studies on post harvest diseases of mango and chemical control of stem end rot and anthracnose. *Ind. Phytopathol.* **47**: 197-200.
264. Sharma, Kavita. (2011). Comparative study of Aeromycoflora in relation to soil mycoflora of Darjeeling tea garden, India. *Recent Research in Science and Technology.* **3**(5): 84- 86

265. Sharma, M.C., Neog, K., Das, A., and Phukan, J.C.D.(2013). Impact of soil fertility and leaf nutrients status on cocoon production of muga silkworm, *Antheraea assamensis* (Helfer) in potential muga growing areas of Assam, India. *International Journal of Current Microb. And Appl. Sc.***2**(9):25-38
266. Sharma, M.S. and Raju, N.S. (2013). Frequency & percentage of occurrence of soil mycoflora in different crop fields at HD Kote of Mysore district. *Int. J. Env. Scs.* **3**(5): 1569-1576
267. Sharma, R. C. and Sharma, Y.P. (1993). Physiochemical alternation of some mulberry leaves infected by blight disease. *Ind. Phytopath.* **43** : 382-384.
268. Sharma, T.C., and Bhattacharjee, K.,(2001). Occurrence of aeromycoflora in banana plants in Kamrup district. *J. Adv. Pl. Sci.* **3** : 21-28.
269. Sharma ,PN., Kaur ,M., Sharma, OP., Sharma, P. and Pathania, A. (2005). Morphological, pathological and molecular variability in *Colletotrichum capsici*, the cause of fruit rot of chillies in the subtropical region of north-western India. *J. Phytopathol.* **153**: 232-237.
270. Sharma Rakesh M.S. and Raju N.S. (2013). Frequency and percentage occurrence of soil mycoflora in different crop fields at HD Kote of Mysore district. *Int. J. Of Env. Sci.* **3**(5): 1569-1575
271. Sharma, S.K., and Gupta, J.S.(1984). Phyllosphere microflora of brown sarson in relation to climatic factors and cultivars. *Ind. Phytopath.* **32**:378-381.
272. Shekhar, MJ., Vinayagamurthy, A., Subramanian, S. and Anbusm, S. (1989). Control of black spot of beer (*Z.maruritiana*). *South Indian Horticul.* **37**:344-345.
273. Shiny, Niharika P., Gaddeyya,G., and Ratna Kumar,P.K. (2013). An investigation on soil mycoflora of different crop fields at Narasannapeta Mandal, Srikakulam district. *Int. Res. J. Environment Sci.* **2**(9):38-44
274. Shivapuri, A. and Gupta, R.B.L. (2002). Evaluation of different fungicides and plant extracts against leaf spot disease of mustard. *Ind. Phytopath.* **54**:372-373.
275. Shree,M.P. and Chandrarama (1999). Biochemical alternation in rust affected leaves tapioca-an important food plant of eri silkworm. *Ind. J. Seric.* **37**: 68-69.
276. Shresth, A.K. and Tiwari, R.D. (2009). Antifungal activity of crude extracts of some medicinal plants against *F.solani* (Mart.) Sacc. *ECOPRINT*.**16**:75-78.
277. Sidhan, G.S. and Bose, S.K. (1980). Epidemiology of angular leaf spot of French bean caused by *Phaeoisariopsis griseola*. *Indian Phytopath.* **33**(1):64-68.

278. Siddiqui, A.A., Babulal, Bhattacharya, A. and Das, P.K. (2000). Nutritional status in morphovariants of som. *Indian Silk*. **21**:18-21
279. Silva, M., Pereira, OL., Braga, IF. and Leli, SM (2008). Leaf and pseudobulb diseases on *Bifrenari harrisoniae* (Orchidaceae) caused by *Phyllosticta capitalensis* in Brazil. *Australasian Plant Disease Notes*. **3**:53-56.
280. Sing, A.; Prakesh, D.M. and Singh, A.B. (1999). Aeromycoflora of grain storage at Udaypur. *Ind. J. Aerobiol.* **12**:15-18.
281. Sing,B. K., Munro,Stracey., Potts,J.M. and Millerd,Peter. (2007). Influence of grass species on rhizosphere microbial community in grass land soil. *Applied Soil Ecol.* **36**:147-153.
282. Singh, J. (1996). Building Mycology-management of decay and health in Buildings, 1<sup>st</sup> ed., Chapman and Hall, London, UK.46-49.
283. Sing, N. I. (1994). Aerobiology and epidemiology of Cercospora leaf spot disease in broad bean (*Vicia Faba* L). *Curr. Trend in Life.*, **20** :171-177.
284. Singh, N.I. and Devi, R.KT. (2001). First report of Broad Been Canker caused by *Pestalotiopsis disseminata* in India. *Pl. Dis.* **85**(2) :229
285. Sing, P. N., Sindhu, I.R. and Singhal, G. (1990). Microfungi associated in non-infected and infected leaves of *Colocasia*. *Acta Botanica India.* **20**: 83-87.
286. Singh, RD. and Prasad, N. (1967). Epidemiological studies of anthracnose of *Dioscorea alata* L. (ratalu)-yam. *Ind. Phytopathol.* **20**: 226-235.
287. Singh, R.S. (2005). Plant disease (8th edition). Oxford and IBH publishing Co. Pvt.Ltd., New Delhi. 439-444.
288. Singh, T.N., Singh, C.K. and Singh, M.S. (2015). Evaluation of different plant extracts for management of powdery mildew of *Quercus serrata* Thunb. Caused by *Phyllactinia corylea* (Pers.) Karst. *Asian Journal of Plant Sci & Res.* **5**(2):77-81.
289. Singh, U.P.; Srivastava,B.P.; Singh, P. K., and Misra, G. D.,(1991). Control of powdery mildews of pea by Ginger extract. *Ind. Phytopath.* **44** : 55-59.
290. Singha A.K., Verma, K.P., Agarwala,K.C., Toorary, N.K. and Thakur M.P.(2004). Antifungal activities of different plant extracts against *Colletotrichum capsici*. *Ad. Plant Sci.***17**(1): 337-338
291. Singha, U. P. S., Prakesh, M., and Singha, A. K. (1992). Variation of chemical constituents in relation to maturity of leaves on mulberry. *Ind. J. Seric.* **32** :196-200.

292. Sinha, S. (1965). Microbiological complex of phyllosphere & disease control. *Indian phytopathology*. **18**: 1-20.
293. Sridhar, T.S. (1978). Some new diseases of fruits. *Current Science* .**47**:71-165.
294. Srinivas, S.G. and Krishnamurthy, Y.L. (2014). Rhizosphere and rhizoplane mycoflora of *Crotalaria pallida* and *Caesalpinia mimosoides*. *NeBIO*.**5**(3):11-14
295. Srivastava, L.S. and Dayal, R. (1986). Studies on rhizosphere mycoflora of *Abelmoschus esculentas*. 14. Antibiosis of Soil microorganisms. *Ind. Phytopath.* **39**:104-106
296. Stepalska, D. and Wolerk, J. (2006) Variation in fungal spore concentrations of selected taxa associated to weather conditions in Cracow, Poland. *Aerobiologia*. **21**: 43-52.
297. Steven, E., Lindow and Johan H. J. (2007). Phyllosphere microbiology. *Current Opinion in Biotechnology*. **22** : 245-151.
298. Sukumar, J., Dandin, S. B. and Bongale, U.D. (1994). Mulberry disease management. Kamatak State Sericulture Research and Development Institute. Publication, Bangalore. 2-3.
299. Sule, I.O. and Oyeyiola, G.P. (2012). Fungal population in the root region of *Cassava* cultivar TMS 30572. *World Journal of Agricultural Sciences*. **8**(1):73-79.
300. Suleiman, M.N. (2011). Antifungal properties of leaf extract of neem and tobacco on three fungal pathogens of tomato (*L. esculentum* Mill). *Adv. in App. Sc. Res.* **2**(4): 217-220
301. Tandon, R.K., Kirtikar and Singh, D.R. (1967). Occurrence of sugarcane rust in Uttar Pradesh. *India Sugar*. **6**:681-683.
302. Takeuchi, Y. (1960). Ability of silkworm (*Bombyx mori* L.) to recover from malnutrition. *The silk news letter*. **5**(8): 6-7
303. Tamini, K.M.; Hadwan, H. A. and Saxena, S.B. (1987). Seasonal influence of rhizosphere fungi of potato. *Ind. Phytopath.* **40** : 555-557.
304. Tewari, S.N. and Nayek, M. (1991). Activity of four plant leaf extracts against three fungal pathogens of rice. *Trop. Agric.* **68**: 373-375.
305. Thakur, M S., Pathok, N. and Saxena, S.B. (1987). Seasonal variation of leaf surface mycoflora of *Azadirachta indica*. *Ind. Phytopath.* **32** :255-256.
306. Thangavelu, K., Chakraborty, A.K., Bhagawati, A.K. and Isa, M. (1998). *Handbook of muga culture*. Bangalore India, CSB.



307. Thangavelu, K. and Sahu, A.K. (1983). Domestication of muga silkworm (*A.assama* WW.); a major breakthrough in muga culture. *Sericologia*.**23**: 153-158.
308. Thimmareddy, H., Prabhuraj,D.K., Bongale U.D., and Dandin, S.B.(1999). Fertility status of mulberry growing soils in mysore seed area, Karnataka. *Indian J. Seric.* **38**(1): 26-29
309. Thangavelu,K., Bhagawati, A.K., and Chakraborty, A.K.(1987). Studies on some wild sericigenous insects of NE India. *Sericologia*. **27**:91-98
310. Thangavelu, K., and Borah,A.(1986). Occurrence of *A.mylitta* Drury (Lepidoptera: Saturniidae) in NE India-distributional significance. *Current Science*. **55**:940.
311. Tikader, A., Vijayan, K., and Saratchandra, B. (2013). Muga silkworm, *A.assamensis* (Lepidoptera: Saturniidae)- an overview of distribution, biology & breeding. *Eur. J. Entomol.* **110**(2): 293-300.
312. Toth, S.J. and Prince, A.L. (1949). Estimation of cation exchange capacity and exchangeable Ca, K and Na content of Soil by flame photometer technique. *Soil Sci.* **67**: 439-445
313. Tripathi, N.N., Sharan, G.S., Kaushik, C.D. and Gupta, I.C. (1987). Magnitude of losses in yield and management of *Alternaria* blight of rapeseed mustard. *Haryana Agri. Univ. Res.* **17**:14-18
314. Umesh,Kumar, N.N.(1992). Biochemical changes in *Shorea robusta* infected by leaf spot disease. *Ind. J. Seric.* **31**: 23-25
315. Unni B.G., Goswami,M., Kakoty,Y., Bhattachajee,M., Wann,S.B., Rajkhowa, G., Das,S., Devi,B.R. and Chutia,A.D. (2009). Indigenous knowledge of silkworm cultivation & its utilization in NER of India. *Ind. J. of Trad. Knowledge.*, **8**(1): 70-74
316. Upadhyaya, M. and Gupta, R.C. (2003). Effect of some medicinal plant extracts on growth of *Curvularia lunata*, a pathogen of *Setaria itali*. *Indian J. Mycol. Pathol.* **20**(2): 145-146.
317. Upadhyaya, J. P. and Pandey, K. K.(2004). Microbial population from rhizosphere and non-rhizosphere soil of pigeon pea : Screening for Resident Antagonist and Mode of Mycoparasitism. *J. Mycol. Pl. Path.* **46** : 7-10.
318. Vala, D.G.(1996). Fungal and bacterial diseases of banana. *Advances in diseases of fruit crops in India.* 38-46.

319. Vankatashwaralu, K. and Monoharachary, C. (1976). Leaf surface mycoflora of some oil yielding and medicinal plants. *GEOBIOS*. **3** :29-30 .
320. Venkataravanappa,V. And Nargund,VB. (2002). Evaluation of different fungicides against *Colletotrichum gloeosporioides* causal organism of anthracnose of mango. Paper presented on Plant disease scenario in southern India. *Annual Meeting Symposium, University of Agriculture Sciences, Bangalore, India*.
321. Vereijssen, J., Schneider, J. H. M., and Jeger, M. J.(2007). Supervised control of *Cercospora* leaf spot in suger beet. *Crop Prot*. **26** : 19-28
322. Verma, R.K., Chaurasia, L., and Katiyar, S. (2008). Potential antifungal plants for controlling building fungi. *Natural Product Radiance*. **7**(4): 374-387
323. Verma, M., Brar, SK., Tyagi, RD., Surampalli, R.Y. and Valero, J.R. (2007). Antagonistic fungi, *Trichoderma* spp.: Panoply of biological control. *Biochem. Engl. J*. **37**:1-20.
324. Vidhyasagar, G. M. and Rajasab, A. H.(2003). Effect of neem, parthenium and garlic extracts on the incidence of powdery mildew of mulberry plant. *Ind. J.Seric*. **40**: 81-85.
325. Vincent, J.M. (1927). Distortion of fungal hyphae in the presence of certain inhibitors. *Nature*.**159**: 180 .
326. Walkley, A. and Black, I. A. (1934). An examination of Degtjareff method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Sci*. **37**: 29-37.
327. Warcup, J. H. (1951). The Ecology of Soil Fungi. *Trans .Br. Mycol. Soc*. **34** :376-400.
328. Warcup, JH.(1957). The ecology of soil fungi. *Transactions of the British Mycological Society*. **34**, 376-379.
329. Wilde, S.A., R.B.Corey, J.G. Iyer and G.K. Viogt.(1985). Soil and plant analysis for tree culture, Oxford and IBH pub. Co. New Delhi.
330. Williams, S.T., Parkinson, D. and Burges, N.A. (1965). An examination of the soil washing technique by its application to several soils. *Plant and Soil*. **22**(2): 167-186
331. Wong, F.P., and Wilcox, WF. (2001).Comparative physical mode of action of azoxystrobin, mancozeb,and metalaxyl against *Plasmospora viticola* (grapevine downy mildew). *Plant disease*. **84**: 649-656.

332. Wood, D.J., Bhatia, I.S., Chakravorty, S., Choudhury, M.N.D., Deb, S.B., Roberts, E.A.H., Yem, E.W. and Willis, A.J. (1954). The estimation of carbohydrates in plant extract by Anthrone. *Biochem. J.* **57**: 508-514.
333. Yadav, 2000. Studies on association between cocoon weight and silk of muga silkworm *Antheraea assama* (Westwood) during different broods. *Int. J. Wild Silk Moths Silk.* **5**: 221-225.
334. Yadav, G.S. and Sampson, M.V. (1987). Muga culture in Mamit Sub division of Mizoram. *Indian Silk.* **26**(5):14-15
335. Yadav, G.S. and Goswami, B.C. (1992). Studies on the foliar constituents of "Som" (*M. bombycina* King) and Soalu (*Litsea polyantha* Juss.). *Sericologia.* **32**:447-451
336. Yadava, G.S. and Goswami, B. (1992). Foliar constituent of som and soalu leaves. *Ind. J. Seric.* **22**: 106-109.
337. Yashodha, H., Hegde, R.K. and Srikant, K. (1993). Physiological studies on *Colletotrichum gloeosporioides* (Penz) Penz and Sacc. causing anthracnose of arecanut. *Karnataka J. Agricul. Sc.* **6**: 411-412.
338. Yemn, E. W. and Willis, A. J. (1954). The estimation of carbohydrates in plant extracts of neem. *Syn. Sci. Seric.* **13** : 165-169
339. Yokoyama, T. (1963). On application of heterosis in Japanese sericulture. In: Proceeding International Genetic Symposium. Suppl. Vol. *Cytology.* 527-531
340. Yokoyama, T. (1996). *Synthesized Science of Sericulture*, Central Silk Board, India. 121-123
341. Zamal, T., Sarmah, B., Hemchandra, O. and Kalita, J. (2010). Global warming and its impact on the productivity of Muga Silkworm (*Antheraea assamensis* Helfer.) *Bioscan.* **1**: 199-209.