

## LIST OF TABLES

Table	Title	Page No
1.	Distribution and acreage of Oak species used as host plant of <i>A. proylei</i> in India	6
2.	States wise Oak tasar raw silk production in India	7
3.	Average leaf yield of per plant of <i>Q. serrata</i> and hectare with application of FYM and NPK in 2013 and 2014 Spring and Autumn season	48
4.	Average Leaf of per plant of <i>Q. serrata</i> in Spring and Autumn season in 2013 and 2014 under different treatments	49
5.	Meterological data records of Oak tasar crop at Umrangso in Spring and Autumn season in 2013 and 2014	51
6.	Rearing performance of <i>A. proylei</i> <b>Jolly.</b> without NPK and FYM application (Control) in 2013 an 2014.	52
7.	Rearing performance of <i>A. proylei</i> <b>Jolly.</b> with application of NPK and FYM in 2013 and 2014	53
8.	Rearing performance of <i>A. proylei</i> <b>Jolly.</b> with application of FYM in 2013 and 2014	54
9.	Effective rate of rearing (ERR) % of <i>A. proylei</i> <b>Jolly.</b> in 2013 2014 under different treatment	55
10.	Fungal isolates of leaf surface of <i>Q. serrata</i> at different status of age in March and April ( Spring season) in 2013 and 2014	58
11.	Fungal isolates of leaf surface of <i>Q. serrata</i> at different status of age in September –October (Autumn season) in 2013 and 2014	60
12.	Relative abundance % Fungal isolates of RS and NRS of ( <i>Quercus serrata</i> ) seedlings during Spring and Autumn season during 2013 and 2014.	63

.13.	Relative abundance % Fungal isolates of RS, NRS and RP soil of <i>Quercus serrata</i> plantation during Spring season 2013 and 2014.	66
14.	Relative abundance % Fungal isolates of RS, NRS and RP soil of <i>Quercus serrata</i> plantation during Autumn season (September-October), 2013 and 2014.	68
15.	Relative abundance % Fungal isolates from air over ( <i>Quercus serrata</i> Plantation during Spring (March-April) and Autumn season (September-October), 2013 and 2014. at 0.75 meter height.	71
16.	Relative abundance % Fungal isolates from air over ( <i>Quercus serrata</i> ) plantation during Spring (March-April) and Autumn season (September-October), 2013 and 2014 at 1.50 meter height.	73
17.	Physio-chemical character of soil under <i>Q. serrata</i> plantation at R.E.C.Umrangso, Farm	75
18.	Foliar constituents of <i>Q. serrata</i> without NPK and FYM application Control in 2013 and 2014	76
19.	Foliar constituents of <i>Q. serrata</i> with application of NPK and FYM in 2013 and 2014	77
20.	Foliar constituents of <i>Q. serrata</i> with application of FYM	78
21.	Leaf constituents of <i>Q. serrata</i> under different treatment in Spring season in 2013(Moisture %, Crude protein %, Crude fibre %,Crude fat%,Ash%,Carbohydrate %) and Statistical analysis	79
22.	Leaf constituents of <i>Q. serrata</i> under different treatments Autumn 2013 and Statistical analysis.	80
23.	Leaf constituents of <i>Q. serrata</i> under different treatments Spring 2014 and Statistical analysis.	81
24..	Leaf constituents of <i>Q. serrata</i> under different treatments Autumn 2014 and Statistical analysis.	82

25.	Reeling parameter of <i>A. proylei</i> <b>Jolly.</b> cocoons under different Treatment	83
26..	Reeling parameter of <i>A.proylei</i> cocoons(Filament length, Denair, Non breakable filament length,Silk recovery%,Reelability %) and Statistical analysis	84